BULLETIN

OF THE

AMERICAN MATHEMATICAL SOCIETY

EDITED BY

EARLE R. HEDRICK

DAVID R. CURTISS WILLIAM R. LONGLEY

WITH THE ASSISTANCE OF

HAROLD T. DAVIS HARRY W. KUHN

THEOPHIL H. HILDEBRANDT DERRICK N. LEHMER

JOHN R. KLINE HENRY L. RIETZ

CAROLINE E. SEELY

VOLUME XL, NUMBER 1

JANUARY, 1934

PUBLISHED BY THE SOCIETY

MENASHA, WIS., AND NEW YORK

1934

PUBLISHED MONTHLY

Whole No. 412 $9.00 a Year

Entered as second class matter April 24, 1926 at the post office at Menasha, Wis., under the act of August 24, 1912.

AMERICAN MATHEMATICAL SOCIETY

Mail Address: 501 West 116th Street, New York, N.Y., for the Secretary, Treasurer, Librarian, and Clerk.

Office: Room 413, Physics Building, Columbia University, New York City.

President: Professor A. B. Coble, University of Illinois, Urbana, Ill.

Secretary: Dean R. G. D. Richardson, Brown University, Providence, R.I.

Associate Secretaries: Professor M. H. Ingraham, University of Wisconsin, Madison, Wis.; Professor J. R. Kline, University of Pennsylvania, Philadelphia, Pa.; Dean T. M. Putnam, University of California, Berkeley, Calif.

Treasurer: Professor G. W. Mullins, Barnard College in Columbia University, New York City.

Librarian: Professor R. C. Archibald, Brown University, Providence, R.I.

Clerk: Dr. Caroline E. Seely, 413 Physics Building, Columbia University, New York City.

Editor-in-Chief of the Bulletin: Professor E. R. Hedrick, University of California at Los Angeles, Los Angeles, Calif.

Managing Editor of the Transactions: Professor R. D. Carmichael, University of Illinois, Urbana, Ill.

Chairman, Colloquium Editorial Committee: Professor Oswald Veblen, University of Texas, Austin, Tex.

Managing Editor of the American Journal of Mathematics: Professor Abraham Cohen, Johns Hopkins University, Baltimore, Md.

Representatives of the Society in the Division of Physical Sciences of the National Research Council: Professors A. B. Coble, E. R. Hedrick, and Dean R. G. D. Richardson.

Representatives on the Council of the American Association for the Advancement of Science: Professors M. H. Ingraham and C. N. Moore.

Chairman, Committee on Arrangements for the Semi-Centennial: Professor T. S. Fiske, Columbia University, New York City.

BULLETIN

OF THE

AMERICAN MATHEMATICAL SOCIETY

EDITED BY

EARLE R. HEDRICK

DAVID R. CURTISS WILLIAM R. LONGLEY

WITH THE ASSISTANCE OF

HAROLD T. DAVIS HARRY W. KUHN
THEOPHIL H. HILDEBRANDT DERRICK N. LEHMER
JOHN R. KLINE HENRY L. RIETZ
CAROLINE E. SEELY

VOLUME XL

JANUARY TO DECEMBER, 1934

PUBLISHED BY THE SOCIETY
MENASHA, WIS., AND NEW YORK

1934

The Collegiate Press
GEORGE BANTA PUBLISHING COMPANY
MENASHA, WISCONSIN

BULLETIN
OF THE
AMERICAN MATHEMATICAL SOCIETY

THE THANKSGIVING MEETING IN PASADENA

The three hundred tenth regular meeting of the Society was held at the California Institute of Technology, Saturday, December 2, 1933. The meeting was called to order at 9:30 A.M. by Professor W. F. Osgood.

The attendance included the following twenty-four members:

O. W. Albert, L. D. Ames, H. M. Bacon, Harry Bateman, E. T. Bell, C. P. Brady, Myrtie Collier, P. H. Daus, Raymond Garver, H. E. Glazier, E. R. Hedrick, Glenn James, D. N. Lehmer, A. D. Michal, W. F. Osgood, T. M. Putnam, G. E. F. Sherwood, D. V. Steed, S. E. Urner, Morgan Ward, L. E. Wear, W. M. Whyburn, C. L. E. Wolfe, E. R. Worthington.

Luncheon for members and their guests was served at the Athenaeum of the California Institute of Technology.

The titles of papers read at the meeting follow. Those whose abstract numbers are followed by the letter *t* were read by title.

1. *On the Nevanlinna and Bosanquet-Linfoot summation methods*, by Dr. A. F. Moursund. (Abstract No. 39-11-325-*t*.)

2. *Ternary continued fractions in a cubic field*, by Professor P. H. Daus. (Abstract No. 39-11-326.)

3. *Ternary continued fractions for cubic units*, by Professor P. H. Daus. (Abstract No. 39-11-327-*t*.)

4. *A census of squares of order* 4, *magic in the rows, columns, and diagonals*, by Professor D. N. Lehmer. (Abstract No. 39-11-328-*t*.)

5. *Matrix differential systems*, by Professor W. M. Whyburn. (Abstract No. 39-11-329.)

6. *On the properties of a determinant function*, by Professor Clifford Bell. (Abstract No. 39-11-330-*t*.)

7. *Interpolation in the mathematics of finance*, by Professor Clifford Bell. (Abstract No. 39-11-331-*t*.)

8. *A set of postulates for "Riemannian" differential geometry in abstract vector spaces*, by Professor A. D. Michal. (Abstract No. 39-11-332.)

9. *A theory of parallel displacement and curvature for "Riemannian" differential geometry in abstract vector spaces*, by Professor A. D. Michal. (Abstract No. 39-11-333-*t*.)

10. *Existence theorems for analytic solutions of ordinary, certain partial, and total differential equations in abstract vector spaces* (preliminary report), by Professor A. D. Michal. (Abstract No. 39-11-334-*t*.)

11. *Analytical and geometrical investigations in normed linear algebras with a finite or denumerably infinite basis* (preliminary report), by Professor A. D. Michal. (Abstract No. 39–11–335-*t*.)

12. *Abstract dynamical systems and contact transformations* (preliminary report), by Professor A. D. Michal. (Abstract No. 39-11-336-*t*.)

13. *Linear connections and non-Riemannian differential geometry in abstract vector spaces*, by Professor A. D. Michal. (Abstract No. 39-11-337-*t*.)

14. *On definitions of abstract polynomials and analytic functions in abstract vector spaces*, by Professor A. D. Michal. (Abstract No. 39-11-338-*t*.)

15. *A critique of the postulate systems for abstract vector spaces and abstract Hilbert space*, by Professor A. D. Michal. (Abstract No. 39-11-339-*t*.)

16. *A partial differential equation connected with the functions of the parabolic cylinder*, by Professor Harry Bateman. (Abstract No. 39-11-340.)

17. *Exponential polynomials*, by Professor E. T. Bell. (Abstract No. 39-11-341.)

18. *On the second case of Fermat's last theorem*, by Professor Glenn James. (Abstract No. 39-11-342.)

19. *On certain methods in dynamics*, by Professor W. F. Osgood. (Abstract No. 39-11-343.)

20. *A law of reciprocity for the entries in a table of linear forms*, by Professor D. N. Lehmer. (Abstract No. 40-1-1.)

21. *The diophantine equation* $X^2 - DY^2 = Z^M$, by Professor Morgan Ward. (Abstract No. 40-1-2.)

22. *A plane rational curve of order* $2k+1$, by Dr. D. C. Duncan. (Abstract No. 40-1-3-*t*.)

23. *The apparent contour of the general* V_3^n *in* S_4, by Professor A. R. Williams. (Abstract No. 40-1-4-*t*.)

T. M. PUTNAM, *Associate Secretary*

THE THANKSGIVING MEETING IN CINCINNATI

The three hundred ninth regular meeting of the Society was held at the University of Cincinnati on Friday and Saturday, December 1–2, 1933. About one hundred persons attended the meeting, among whom were the following fifty-one members of the Society:

W. E. Anderson, H. F. Archibald, W. L. Ayres, F. R. Bamforth, G. M. Bareis, I. A. Barnett, Henry Blumberg, Louis Brand, R. S. Burington, R. D. Carmichael, A. B. Coble, L. W. Cohen, L. L. Dines, H. H. Downing, J. D. Elder, Arnold Emch, L. M. Graves, Harris Hancock, E. H. Hanson, R. E. Hundley, W. R. Hutcherson, M. H. Ingraham, H. S. Kaltenborn, H. J. Kersten, J. H. Kindle, Lincoln LaPaz, C. G. Latimer, F. E. LeStourgeon, C. I. Lubin, C. C. MacDuffee, C. W. Mendel, G. M. Merriman, H. A. Meyer, C. N. Moore, M. E. Mullings, Alta Odoms, S. E. Pence, H. S. Pollard, Tibor Radó, G. Y. Rainich, S. E. Rasor, F. W. Reed, C. E. Rhodes, G. W. Starcher, Guy Stevenson, A. H. Tappan, H. P. Thielman, Sister M. D. Thuener, O. Veblen, K. P. Williams, E. I. Yowell.

On Friday afternoon Professor R. D. Carmichael gave an address entitled *Functions of exponential type.* On Saturday afternoon there was a joint meeting with the American Physical Society on *Spinor analysis* at which Professor Otto Laporte spoke on *Spinors and their significance in modern physics* and Professor Oswald Veblen spoke on *Spinor analysis.* There was also a large amount of discussion at this meeting.

Friday evening a dinner was held at the Hotel Sinton-St. Nicholas. About fifty persons attended thedinner. Professor Harris Hancock of the University of Cincinnati welcomed the visiting mathematicians, and President Coble responded to his welcome expressing the pleasure of the group in meeting at Cincinnati.

Saturday noon the mathematicians and physicists met at a joint luncheon at the Mecklenburg Restaurant. About seventy-five persons attended this luncheon.

President A. B. Coble and Professor C. N. Moore presided at the meetings on Friday afternoon, President Coble and Professor L. L. Dines on Saturday morning, and Professor J. H. Van Vleck at the joint session Saturday afternoon.

A very pleasant feature of the meeting was the tea served by the ladies of the University of Cincinnati to the visiting mathematicians and physicists on Friday afternoon.

The titles of papers read at the meeting follow. Those whose abstract numbers are followed by *t* were read by title. Mr. P. M. Pepper was introduced by Professor C. N. Moore, Mr. H. E. Vaughan by Professor W. L. Ayres, Miss Marjorie Leffler by Professor Tibor Radó, Mr. W. C. Mitchell by Professor C. N. Moore, and Mr. E. S. Akeley by Professor Cornelius Lanczos.

1. *A classification of plane cubic curves under the affine group by means of arithmetic invariants*, by Professor R. S. Burington. (Abstract No. 40-1-5.)

2. *Some remarkable sextic space curves*, by Professor Arnold Emch. (Abstract No. 39-11-301.)

3. *Affine-polygenic functions*, by Mr. P. M. Pepper. (Abstract No. 40-1-6.)

4. *A significant vector in the theory of surfaces*, by Professor Louis Brand. (Abstract No. 40-1-7.)

5. *On the units in a cyclic field*. by Professor C. G. Latimer. (Abstract No. 40-1-8.)

6. *Some theorems on connected sets*, by Mr. H. E. Vaughan. (Abstract No. 40-1-9.)

7. *On the relation between local connectivity and another property*, by Professor W. L. Ayres. (Abstract No. 40-1-10.)

8. *On the relative stability of the median and arithmetic mean*, by Professor H. S. Pollard. (Abstract No. 40-1-11.)

9. *The first and second variations of an n-tuple integral in the case of variable limits*, by Professor H. A. Simmons. (Abstract No. 39-11-303-*t*.)

10. *A determination of all normal division algebras of degree* 4 *over F of characteristic* 2, by Professor A. A. Albert. (Abstract No. 39-11-300-*t*.)

11. *On certain imprimitive fields of degree* p^2 *over P of characteristic p*, by Professor A. A. Albert. (Abstract No. 39-11-299-*t*.)

12. *Normal division algebras over a modular field*, by Professor A. A. Albert. (Abstract No. 39-11-302-*t*.)

13. *The flexure problem for rectangular beams with slits*, by Professor D. F. Gunder. (Abstract No. 39-11-298-*t*.)

14. *An existence theorem for generalized pfaffian systems*, by Professor J. M. Thomas. (Abstract No. 39-11-297-*t*.)

15. *Note on potential theory in n-space*, by Professor J. J. L. Hinrichsen. (Abstract No. 39-11-321-*t*.)

16. *On measures of contingency* (preliminary report), by Professor F. M. Weida. (Abstract No. 39-11-324-*t*.)

17. *On a problem of Knaster and Zarankiewicz*, by Professor J. H. Roberts. (Abstract No. 39-11-323-*t*.)

18. *Algebraic functions of Stepanoff almost periodic*, by Dr. R. H. Cameron (National Research Fellow). (Abstract No. 40-1-12-*t*.)

19. *On a lemma of Fejér*, by Professor Lincoln LaPaz. (Abstract No. 40-1-13.)

20. *A lemma in potential theory* (preliminary report), by Miss Marjorie Leffler. (Abstract No. 39-11-322.)

21. *Singularities on maps of surfaces*, by Professor G. Y. Rainich. (Abstract No. 40-1-14.)

22. *Transformation of an integro-differential equation by means of a Volterra transformation*, by Professor I. A. Barnett. (Abstract No. 40-1-15.)

23. *Some properties of Stieltjes integrals*, by Mr. H. S. Kaltenborn. (Abstract No. 40-1-16.)

24. *On convergence factors for double series that are summable of non-integral orders*, by Professor C. N. Moore. (Abstract No. 40-1-17.)

25. *Borel-Sannia summability for double series*, by Mr. W. C. Mitchell. (Abstract No. 39-11-320.)

26. *On the Cesàro mean of double Fourier series*, by Miss Alta Odoms. (Abstract No. 40-1-18.)

27. *An invariantive characterization of quadratic differential forms*, by Dr. E. S. Akeley. (Abstract No. 40-1-19.)

28. *A new proof of a theorem of Denjoy, Young, and Saks*, by Mr. E. H. Hanson. (Abstract No. 40-1-20.)

29. *The equivalence of n-tuples of matrices* (preliminary report), by Professor O. E. Brown. (Abstract No. 40-1-21-*t*.)

M. H. INGRAHAM, *Associate Secretary*

BIRKHOFF ON AESTHETIC MEASURE

Aesthetic Measure. By George D. Birkhoff. Cambridge, Harvard University Press, 1933. xvi+226 pp., 23 plates, 10 in color. $7.50.

Every reader of this Bulletin will find this work thought-provoking. Its first appeal is to the analytically-minded aesthetician; but there are questions and implications far beyond any specialist's domain. The reviewer will therefore examine *Aesthetic Measure,* firstly, as to how well it solves its self-imposed problem, and, secondly, as to the relevance of its contribution to aesthetics, this term being used in its broadest sense.

In Chapter 1, Professor Birkhoff presents the following mathematical formulation of the fundamental problem: "Within each class of aesthetic objects, to define the order O and the complexity C so that their ratio $M=O/C$ yields the aesthetic measure of any object in the class." In subsequent chapters, this problem is solved for various simple classes of aesthetic objects in great detail. The author distinguishes between "formal" and "connotative" associations, and explains that "our attention will be directed almost exclusively toward the formal side of art, to which alone the basic formula of aesthetic measure can be quantitatively applied," but with "no intention of denying the transcendent importance of the connotative side in all creative art."

In Chapters 2, 3, and 4, the author elaborately applies his theory to polygonal forms, ornaments and tilings, and vases—that is, to the more elementary fields. The reader stands amazed at the author's thoroughness. Here minor criticisms are beside the point; for the author does seem to establish the validity of his formula. In Chapter 2, there is recorded the aesthetic measure of 90 polygonal forms, pictured in colored plates. In Chapter 3, five interesting full-page plates illustrate species of one- and two-dimensional ornaments. But the discussion preliminary to application of the theory, though a neat piece of analysis, is sufficiently mathematical to become formidable reading. In Chapter 4, we meet a difficulty. That the theory "here is to be regarded as more questionable than the theories of polygons and ornaments" suggests that the more abstract vase form has higher expressiveness. And we sense trouble ahead in dealing with an art so abstract as music. The author himself concedes, in closing the chapter on vases: "It is an interesting question as to how the explicit knowledge that these simple relations exist in a given case affects the aesthetic judgment. In my opinion, the effect is slightly adverse, for, in all fields of art, it is the intuitively felt relationships which are the most enjoyed."

In Chapters 5, 6, and 7, we come to music. In these chapters on chords, harmony, and melody, we recognize the author's perseverance in acquainting himself with the many technical matters involved and his patience in enumeration. We gain the impression that the author, willing to admit the "almost transcendental expressive power" of music and its "deep and almost universal appeal," has yet a tendency to pass lightly over certain features which, although irrelevant for him, are, for many, vital. There is a wide-spread conviction that an art-work is significant because of what it "means" rather than what it "says." We hold, with Birkhoff, that Gurney's point of view—that

music is Ideal Motion, whose "essential characteristic . . . is an absolutely unique beauty perceived by an absolutely unique faculty . . . "— is hardly tenable. Yet must a science of aesthetics necessarily be analytical in the sense of Birkhoff to be other than mystical?

The chapters on harmony and melody present, it is true, a valid formula. But what, precisely, is its significance for aesthetics? We find merely a rationalization of rules previously discerned through a study of the masters—and made conveniently accessible by Prout's *Harmony* and *Counterpoint*. It is only natural that the theory should turn out to be both necessary and sufficient: the author finds that no good melody can have a low M, and that no poor melody can have a high M.

Now to some aestheticians, formal structure explains the whole power of music. For the musician, however, the major emphasis is elsewhere. The musically receptive and discriminative listener requires knowledge neither of musical theory nor of aesthetic measure. His query as to the formal is merely: shall it be subjected to rational or to intuitive analysis?

In these chapters, it would not be difficult for the theoretician to propound some interesting questions. For example, on page 148, in assigning $M=15$ to the fifth chord sequence in the first brace, and $M=16$ to the eighth in the second brace, does not the author overlook vital elements in musical significance? And how about rests? Every reader of musical material, from either the creative or the analytical side, has heard that "it is easy to write notes but hard to write rests"—a remark generally attributed to Mozart. Any analysis which does not take into account the soul-stirring effect of measures of rest in the first movements of Beethoven's Fifth Symphony and Mozart's G Minor Symphony, can hardly escape classification as inadequate even from an elementary standpoint.

Birkhoff's analysis of the formal factors in poetry (Chapter 8) renews interest in the ever-burning question as to the relative merits of views held by various schools. Birkhoff says that "the fundamental aim is always to achieve the terse, imaginative expression of the poetic idea in metric form by use of language of unusual musical quality." In contrast with this notion stands Professor Housman's* definition of poetry: "I think that to transfuse emotion —not to transmit thought, but to set up in the reader's sense a vibration corresponding to what was felt by the writer—is the peculiar function of poetry." Housman considers poetry "more physical than intellectual," but maintains that "the majority of mankind notoriously and indisputably do not . . . possess the organ by which poetry is perceived." In short, although poetry is not so abstract and powerful a medium of expression as music, the vital aim is that of music: to stir our feelings through voicing them; and the crucial test is how deeply we are stirred! Moreover, in poetry, as in music, the highest desideratum is that perfect unification of form and content for which all art strives; and if we are concerned with what poetry is rather than with what it is about, it is doubtful how much any analysis, of factors abstractly separated, can contribute to our understanding of the fused whole, where the synthesis is so subtle as virtually to defy analysis. We are driven again to query whether the

* A. E. Housman, *The Name and the Nature of Poetry*, Macmillan, 1933.

meaning of art is so much the information it conveys as what *it* imparts by way of power to apprehend and evaluate experience.

The author's appraisals are found in the two concluding chapters, on *Earlier Aesthetic Theories* and on *Art and Aesthetics*. The contributions of earlier writers, beginning with Plato and Aristotle, are interpreted in terms of the quantitative theory. To the author "it seems almost obvious that aesthetics, if it is to be scientific, must be approached from the analytic point of view and must concern itself chiefly with the formal aspects of art." Thus, when Birkhoff comes to Croce, for whom art is the "expression of impressions" and "lyrical intuition," he observes: "Such general philosophical definitions and classifications, however true, can never serve as the point of departure for a science of aesthetics. They are self-limited and form a kind of philosophic citadel from which an attack upon any and all more definite conclusions can be conveniently made."

Now the science of aesthetics envisaged by Vico, and later brought to flower by Croce, excludes by its very nature the type of analysis favored by Birkhoff. But this does not mean that Vico or Croce have offered us a mystical theory, or something vague in outline or content. To the reviewer it has always seemed that Croce, in his *Aesthetic*, penetrates toward the real heart of the matter. He argues ably for a science of intuitive knowledge comparable in importance to a science of intellectual knowledge, and stresses the fact that: "The difference between a scientific work and a work of art, that is, between an intellectual and an intuitive fact, lies in the difference of the total effect aimed at by their respective authors. This it is that determines and rules over the several parts of each, not these parts separated and considered abstractly in themselves."

In this chapter, the author also quotes, in part, two paragraphs from the end of Helmholtz's monumental work on *Sensations of Tone*. Birkhoff, although he finds no indication of the quantitative outlook, concludes that Helmholtz adheres to the general analytic point of view; for the latter says: "No doubt is now entertained that beauty is subject to laws and rules dependent on the nature of human intelligence . . . " But, as we read the chapter from which Birkhoff quotes, it seems to us that Helmholtz distinctly subordinates the analytic point of view; indeed, he says: "It is precisely from that part of its regular subjection to reason which escapes our conscious apprehension that a work of art exalts and delights us, and that the chief effects of the artistically beautiful proceed, *not* from the part which we are able fully to analyze." And Helmholtz decides not "to proceed further into the esthetics of music . . . The real difficulty would lie in the development of the psychical motives which here assert themselves. Certainly this is the point where the most interesting part of musical esthetics begins, the aim being to explain the wonders of great works of art, and to learn the utterances and actions of the various affections of the mind."

In the concluding chapter, Birkhoff indicates very briefly how application of his theory may be made, in decorative design, painting, sculpture, architecture, and music, to include the qualitative aspects of form. It is shown that certain well known maxims—embodied in the principle of "unity in variety"—are contained implicitly in the basic formula. But if these maxims have been the point of departure for formalism ever since antiquity, does not the author

here expose himself to the criticisms levelled at the formalists? After some remarks on evolution of art and on creative art, the chapter, and therewith the book, closes with a description—far from convincing to us—of the services which the author believes his theory capable of performing both in the aesthetic experience and in the creative process.

As we lay aside *Aesthetic Measure*, our first thought is that, for an estimate of its relevance, we should view it against a sufficiently broad philosophic background. The complete picture must embrace not only man's intuitive and conceptual activities, but also his practical activities, economic and ethical.

We affirm that the intuitive and the conceptual are indisputably distinct. Yet may there not be some parity connecting such representative theoretical activities as music and mathematics? If found, this parity should throw into even sharper relief the difference between the intuitive and the conceptual—by revealing to us more clearly those aspects in which the two are not comparable.

The reviewer's contention is that the parity we seek is to be found in the striking parallelism between, on the one hand, the manner in which mathematics, through correspondence, "functions" with formal or quantitative aspects of the concrete to extend our range of conceptual knowledge, and, on the other hand, the manner in which music, through association, "functions" with connotative or affective aspects of the concrete to enlarge our sphere of intuitive knowledge. Here "function" refers to the interplay between the real and the ideal. Thus, on Dewey's view, both art and science would appear as controls for progress; to Santayana, both music and mathematics would glory in being wholly free and yet useful. We cannot pass, in the sense of establishing identical correspondences, from any one realm to another; yet, through comparison and suggestion, we are able to obtain ever-increasing knowledge of ourselves and of our physical surroundings. We dare to opine that this functional dependence of progress on activities both theoretical and practical has virtually the same characteristics whether in the domain of the intuitive or the conceptual. It must be left to the reader to realize the parity we have conjectured and the force of its implications. Our own inference from the generalization just advanced is that, until we totally separate music and mathematics, art and science, and proceed on the basis of their parity, we shall not be making headway.

C. A. GARABEDIAN

PAPPUS'S COLLECTION

Pappus d'Alexandrie. La Collection Mathématique. Œuvre traduite pour la première fois du Grec en Français, avec une introduction et des notes. By Paul Ver Eecke. Paris and Bruges, Desclée De Brouwer, 1933. 2 vols. 128+364+507 pp.

It seems strange that such a notable classic as the *Collection* of Pappus should never have been completely translated into any modern European language until M. Ver Eecke undertook the task of preparing it for this French edition. Commandino had put it in Latin in 1588, this being the best way of making it known to scholars of that time; but even this translation, elaborate as it was, has since been shown to be quite imperfect. Three centuries had then to elapse before any attempt was made at a definitive edition of the work. This edition was the work of Dr. Friedrich Hultsch, a scholar of highest rank, and it presented both the Greek and the Latin versions based upon a critical study of the earliest and most complete manuscripts extant. Sir Thomas Heath speaks of it as "one of the first monuments of the revived study of the history of Greek mathematics in the last half of the nineteenth century." Upon this edition M. Ver Eecke bases his translation. It should be said, however, that there were numerous partial translations, such as that made by Dr. Gerhardt, of Books VII and VIII (Halle, 1871).

Since the publication of the Commandino edition the date of Pappus has been more clearly fixed, and we now know that he flourished in the reign of Diocletian (284–305). Out of his many works, most of which are lost, his *Mathematical Collection* (*Mathematikai synagogai*) is the best known. It is a kind of synthesis of mathematics as known in his time, with many historical notes, and is the source of much of our knowledge of the works of the classical period.

In an elaborate Introduction of more than a hundred pages M. Ver Eecke gives a general survey of the known facts relating to Pappus, a statement of the nature of each of the extant Books of the *Collection*, and a list of the various editions consulted. The great value of the translation lies not only in its apparently careful rendering of the Greek in French, but in the footnotes. The latter contain much information concerning the work itself, together with a large number of proofs in modern form and of bibliographical notes.

The contents of the several Books, beginning with the fragment of Proposition 14 of Book II, are so fully stated by Sir Thomas Heath in his *History of Greek Mathematics* (vol. II, pp. 361–439) as to call for little further notice of the subject. It should be said, however, that the arrangement followed by Pappus differed greatly from that of Euclid or of Appollonius. The work is rather, as the title indicates, a collection of notes on geometry, the number theory, and the geometric algebra of the Greeks, than a textbook on any one of these subjects. Of the several divisions, Book IV is of particular interest because of Pappus's treatment of the generalized Pythagorean Theorem; of the *arbelos* (shoemaker's knife) with its inscribed circles; of the spirals studied by Euclid, Archimedes, and Conon; of "a certain line used by Nicomedes for

the duplication of the cube" (the conchoid); and of the *tetragonizousa gramma*, which in the Latin of Commandino appears as the *linea quadrans*, known at the present time by the late Latin term *quadratrix* (French *quadratrice*). Book V is also particularly interesting because of its geometric content, especially as relates to isoperimetry; maximum and minimum lines, surfaces, and solids; the treatment of the sphere and cylinder by Archimedes; and the regular and semi-regular solids. Book VII, the *Analuomenos topos* (Treasury of Analysis), gives the clearest statement of the Greeks relating to the distinction between analysis and synthesis. It also contains the theorem relating to the volume generated by the revolution of a plane surface about an axis, the so-called Guldin Theorem. Book VIII treats of theoretical mechanics, and Sir Thomas Heath mentions the fact that Carpus (*fl.* about 100 A.D.), to whom Pappus here refers, "seems in reality to have been anticipating the modern view of an angle as representing divergence rather than distance" and to have had the idea of rotational as distinct from linear measure.

The French language has been the medium used by two noteworthy historians of mathematics in the present century, Paul Tannery (most of whose works have been edited posthumously by or under the direction of his widow) and M. Ver Eecke. The latter has given us other excellent translations of the Greek classics,—the works of Archimedes, Apollonius, Diophantus, Theodosius, and Serenus,—a remarkable achievement in only a few of the years allotted to man. Neither of these historians was a member of a university faculty; neither was closely connected with any of his country's schools; Tannery held a post in the French tobacco monopoly, and Ver Eecke is an "Ingénieur des mines" in Belgium and "Inspecteur général honoraire du travail." Each carried on the traditions of such laymen as Vieta, Fermat, Descartes, and Leibniz—great leaders in the discoveries of knowledge and in opening new vistas to all "who have eyes to see."

One regret will be voiced by all scholars who have occasion to read these volumes, the publication of which has been made possible by the Fondation Universitaire de Belgique—they have no index either to each part or to the two together. There is not even a table of contents. Without such aids the books lose much of their value. A reader who wishes to find a reference to Guldin, the quadratrix, or the glossocome, for example, must thumb through more than 800 pages, and even then will be fortunate if he finds what he seeks. It is no spirit of fault-finding criticism which will prompt scholars generally to express regret that such an omission should be allowed to detract so much from the value of a work of this importance.

That this and the other translations of M. Ver Eecke, mentioned above, should be in every college or university library need hardly be suggested. They are invaluable to all students of the history of Greek culture in general and mathematics, physics, and astronomy in particular.

DAVID EUGENE SMITH

BANACH ON LINEAR OPERATIONS

Théorie des Opérations Linéaires. By Stefan Banach. Warsaw, Monografje Matematyczne, Vol. I, 1932. vii+254 pp.

The present book, the first of the series published by the Monografje Matematyzcne, is an enlarged and considerably modified translation of the *Teorja Operacyi* (in Polish) by Banach. It represents a noteworthy climax of long series of researches started by Volterra, Fredholm, Hilbert, Hadamard, Fréchet, F. Riesz, and successfully continued by Steinhaus, Banach, and their pupils. In a short review it is impossible to give even an approximate idea of the richness and importance of the material, entirely new to a large extent, which is gathered in this book. Roughly speaking the book treats of the field which was touched upon by Hildebrandt in his excellent address on linear functional transformations in general spaces,* but naturally is considerably more extensive and intensive. The theory of linear operations is a fascinating field in itself but its importance is still more emphasized by numerous beautiful applications to various problems of analysis and function theory (real and complex). A number of these applications are contained in the book; a still larger number of them, increasing from year to year, are being inspired by the ideas developed in the book. An attentive reader will find a constant source of inspiration in this monograph which undoubtedly will exercise a great influence on the further development of the subject. A description of the contents follows.

In the Introduction (19 pp.) the author recalls some basic properties of Lebesgue and Stieltjes integrals, and of metric spaces. Particular attention is given to subsets of such spaces, which either are Borel measurable or possess the property of Baire, and also to Borel measurable operations transforming one metric space into another. In Chapter 1 (*Groupes*, 6 pp.) the author specializes the metric space under consideration by adding the usual postulates for groups (with respect to the operation "+", with inverse designated by "−") and also the two postulates concerning limits:

$$x_n \to x \quad \text{implies} \quad (-x_n) \to (-x);$$

$$x_n \to x \text{ and } y_n \to y \quad \text{implies} \quad (x_n + y_n) \to (x + y).$$

It is proved that every subgroup which possesses Baire's property and is of the second category coincides necessarily with the whole space (assumed to be complete). The additive and linear (additive+continuous) operations are introduced at this early stage. It is proved that the set where a sequence $\{U_n(x)\}$ of linear operations converges, or even where it is only bounded, is either the whole space or a set of the first category (the space is assumed to be connected). Chapter 2 (*Espaces vectoriels généraux*, 9 pp.) deals with linear spaces and additive operations defined on such spaces. The central point here is an important theorem concerning a possibility of extending to the whole space

* This Bulletin, vol. 37 (1931), pp. 185–212.

an additive functional (a real-valued operation) defined on a linear sub-space. Theorems of this kind originated with F. Riesz (in the case of space of continuous functions) and the theorem in question was proved independently by Hahn. As an application it is shown that every bounded sequence has a "limit" which is distributive and possesses some other properties of the usual limit, and is always enclosed between the limits of indetermination of the sequence. An analogous application is made to show that an arbitrary bounded function may be supplied with an "integral," as well as every bounded linear set with a "measure." In Chapter 3 (*Espaces du type* (*F*), 18 pp.) a metric complete space is specialized to be linear and to satisfy three additional postulates:

$(x, y) = (x - y, 0)$;

$h_n \to 0$ implies $h_n x \to 0$, where $\{h_n\}$ is a sequence of constants;

$x_n \to 0$ implies $hx_n \to 0$, where h is a constant.

A fundamental property of linear operations (transforming such a space E of type (*F*) into a set of another space E_1, also of type (*F*)) is that the range of a linear operation is either a set of the first category in E_1 or E_1 itself, while, in the case when the transformation (linear) is one to one, it is also bicontinuous. This chapter contains important applications to the theory of continuous non-differentiable functions, the continuity of solutions of partial differential equations, and the solution of certain systems of infinitely many equations of a general type.

The next specialization is furnished by *normed* spaces which are metric and linear with the metric $\|x\|$ satisfying the postulates

$$\|x + y\| \leqq \|x\| + \|y\|, \quad \|cx\| = |c| \, \|x\|,$$

where c is a constant. A normed space, if it is complete, is designated as a space (*B*). Such spaces have been also introduced by Wiener, independently and almost simultaneously with Banach. Chapter 4 (*Espaces normés*, 25 pp.) deals with normed spaces in general. Here we find a determination of the form of the most general linear functional in various important special spaces, such as spaces of all convergent sequences, (*c*); of sequences $x = \{\xi_n\}$ with

$$\sum |\xi_n|^p < \infty, \quad (l^p);$$

of continuous functions, (*C*); of measurable functions $x(t)$ with

$$\int_0^1 |x(t)|^p dt < \infty, \quad (L^p).$$

We find also interesting applications to the theory of approximation (in various spaces) to a given function by means of linear combinations of functions of a given set, and to the moment problem. Chapter 5 (*Espaces du type* (*B*), 18 pp.) deals especially with normed complete spaces. Here we find the fundamental result that if $\{U_n(x)\}$ is a sequence of linear operations, bounded at each point of the second category of a space E of type (*B*), then the sequence of norms $\{\|U_n\|\}$ is bounded. This theorem is the most general among similar results of more or less special nature obtained by various other authors. Some interesting applications to the comparative study of definitions of summability are also found in this chapter. We mention also some results of importance for the theory of Fredholm integral equations with kernels of "bounded"

type, which are out of the reach of the more powerful theory developed by von Neumann and Stone. Chapter 6 (*Opérations totalement continues et associées*, 10 pp.) introduces the notions of the adjoint (or conjugate) operation, and of a completely continuous operation, whose importance is so well known in the more special theories of integral equations and bilinear forms. Chapter 7 (*Suites biorthogonales*, 9 pp.) contains various applications to the general theory of biorthogonal expansions. Of a particular elegance are the results concerning factor-sequences leaving invariant the "class" (L^p) or (C) of a Fourier series. Chapter 8 (*Fonctionnelles linéaires dans les espaces du type* (B), 18 pp.) is devoted to a more detailed study of various sets of linear functionals in spaces (B). Three important notions are introduced and discussed, those of sets of functionals which are regularly closed, transfinitely closed, and weakly closed. The two first are shown to be equivalent for every space (B), while all three are equivalent in case the space (B) is also separable. Conditions are given in order that a given set of functionals coincide with the space of all the functionals (the conjugate space). Chapter 9 (*Suites faiblement convergentes d'éléments*, 12 pp.) gives a discussion of weak convergence, an important notion introduced in special cases by F. Riesz. After all these preparations the author can easily develop his beautiful theory of linear functional equations in Chapter 10 (*Équations fonctionnelles linéaires*, 20 pp.). As very special instances of applications of results of this chapter we may mention the theory of non-singular, and some singular, Fredholm integral equations and of Hilbert's theory of limited and completely continuous bilinear forms. The power of the method (the basic ideas of which were previously introduced by F. Riesz and by Hildebrandt) is due to the fact that it is free from any special way of representation of operations in question, such as integrals or matrices, but deals directly with operations themselves.

The last two chapters treat of entirely different questions. These chapters, as well as the Appendix at the end of the book, were not present in the Polish edition of the book. Chapter 11 (*Isométrie, equivalence, isomorphie*, 28 pp.) discusses various types of correspondences (all of them one to one) between two spaces E and E_1 of type (B). These spaces are called isometric if the correspondence preserves the norm, isomorphic if this correspondence is linear, and equivalent if they are isomorphic and isometric. Some of these notions can be of course extended to more general spaces. A rotation of E about a point x_0 is defined as an isometric transformation of E into itself, which leaves invariant the point x_0. Various beautiful results are found in this chapter, of which we mention only three: Every isometric transformation of a linear normed space into another space of the same type is linear, provided it transforms the zero element into the zero element (Mazur and Ulam). Let Q and Q_1 be two metric complete and compact sets. A necessary and sufficient condition that they be homeomorphic is that the spaces (C_Q) and (C_{Q_1}) of real-valued continuous functions defined over Q and Q_1 respectively be isometric. Every separable space of type (B) is equivalent to a closed linear sub-space of the space (C) of continuous functions; every metric separable space is isometric with a sub-space of (C). The expressions for the most general rotations in spaces (C), (c), (L^p), (l^p) are also found in this chapter. Chapter 12 (*Dimensions linéaire*, 15 pp.) introduces a method of comparing various spaces according to their *lin-*

ear dimensions. We say that, for two spaces of type (F), (1) $\dim_l E \leqq \dim_l E_1$ if E is isomorphic with a linear closed sub-space of E_1; that $\dim_l E = \dim_l E_1$ if also (2) $\dim_l E_1 \leqq \dim_l E$, while $\dim_l E < \dim_l E_1$ if (1) is satisfied but (2) is not. If neither (1) nor (2) is satisfied, the linear dimensions of E and E_1 are said to be incomparable. It should be observed that, while two isomorphic spaces are of the same linear dimensions, there exist separable spaces (B) which are not isomorphic although their linear dimensions are the same. Special attention is given here to spaces (L^p) and (l^p). It is shown that if $\dim_l(L^p) = \dim_l(l^q)$, where $p, q > 1$, then $p = q$; if $1 < p < 2 < q$, then the linear dimensions of (L^p) and (l^q) are incomparable; if $1 < p \neq 2$, then $\dim_l(L^2) < \dim_l(L^p)$; if $\dim_l(L^p) < \dim_l(L^q)$, where $p, q > 1$, then $p = q = 2$; the condition $p = q = 2$ is necessary and sufficient that $\dim_l(L^p) = \dim_l(l^q)$; if $1 < p \neq 2$, then $\dim_l(L^p) > \dim_l(l^p)$. The Appendix (18 pp.) contains an additional discussion of the weak convergence of elements and of functionals. The book closes with 21 pages of additional remarks containing historical information and statements of various generalizations of the results in the text, as well as numerous unsolved problems. These remarks reveal that the whole subject is in a state of vigorous development. The usefulness of many of these remarks would be considerably greater if they contained hints indicating how the results in question have been obtained. The exposition as a rule is clear and detailed, although not easy in several places. The number of misprints (of which only two are corrected in the list of Errata) is not negligible.

In conclusion the reviewer may express his conviction that Banach's monograph should occupy a permanent and honorable place on the desk of every one who is interested in the theory of linear operations, to be replaced only by subsequent, corrected and augmented editions, which undoubtedly will follow before long.

J. D. TAMARKIN

SAKS ON INTEGRATION

Théorie de l'Intégrale. By Stanislaw Saks. Warsaw, Monografje Matematyczne, Vol. II, 1933. vii+290 pp.

The present volume is the second one of the series Monografje Matematyczne. It is a translation, entirely revised and augmented by several important chapters, of the author's Polish book, *Zarys Teorji Calki*, Warsaw, 1930. It fills in a serious gap in the literature of the real function theory, and of the theory of differentiation and integration, which has been acutely felt during all the recent period of vigorous growth and development of these disciplines. While Hermite, together with a large part of his contemporaries and immediate successors, including Poincaré, contemplated with horror the pathological cases of functions without derivatives, the ideas and methods created precisely for handling such bad cases turned out to be extremely fruitful and indispensable for the treatment of most classical and venerated problems of analysis. Far from bringing in the feared anarchy and disorder, they have allowed us in many instances to reach a harmony and completeness of results which were entirely out of the reach of older classical methods. The problems

which require essentially a fundamental knowledge of the most modern chapters of the real function theory are "everywhere dense" in the field of analysis, and the mathematician who wants to make any headway in the theory of functions of a complex variable, of differential and integral equations, of Fourier series, and the like, will find himself badly handicapped if he is not well equipped on the side of the real function theory, and particularly of the theory of differentiation and integration. But these are precisely the chapters which so far have been lacking an adequate treatment in the literature. Except for the treatise of Hobson, which still holds its honorable place, we are unable to name any other book or monograph where a serious attempt has been made to unify the numberless results scattered throughout all mathematical periodicals and to give a general picture of the present state of the subject. Such an attempt, and a fully successful one, is represented by the book of Saks. Its main characteristic unifying feature is the systematic use, from the very beginning, of the notion of additive functions of sets. This allows him to bring in, with greatest clearness and economy of thought and notation, the *descriptive* theory of various integrals he discusses. At the same time it leads very naturally to fundamental problems of the modern differentiation theory, and to an extensive use of Dini's derived numbers and of approximate derivatives and derived numbers, as well as to the question of the unique determination of an additive function of sets by means of its derived numbers of various sorts. The systematic development of these notions constitutes the very nerve of the book. A mathematician who will not avail himself of the rich source of information brilliantly presented in Saks' book will deprive himself of the use of valuable ideas and tools which would be of help to him during all his "analytical" career. In publishing the present monograph and the preceding one by Banach, the Monografje Matematyczne have set up a high standard which will be difficult to match. We congratulate them on this happy start and we wish to attract to this important undertaking all attention of the mathematical world, which it justly deserves. The next books which are announced in this series are: K. Kuratowski, *Topologie* I (Vol. 3); W. Sierpinski, *Hypothèse du Continu* (Vol. 4), and A. Zygmund, *Trigonometric series* (Vol. 5).

The first five chapters of Saks's book treat of the theory of functions of bounded variation, Lebesgue measure and Lebesgue integral, together with a short discussion of the Riemann-Darboux-Stieltjes integral. Without losing anything in simplicity or elegance but gaining in generality and clearness, the author discusses functions of several variables, rather than those of a single variable. An excellent example of application of the foregoing theory presents itself in the theory of areas of surfaces given in explicit form $z=f(x, y)$, as developed by Lebesgue, Geöcze, Tonelli, and Radó. The exposition of this beautiful theory has never been given before in any book; it occupies Chapter 6. Chapter 7 deals with the notion of a Perron integral and its relationship with the integral of Lebesgue. A generalization of the majorant and minorant functions introduced by La Vallée Poussin plays an important role here. Chapters 8 (*Functions of generalized bounded variation*) and 9 (*Theorems on derived numbers*) contain material which is important in itself and at the same time prepares the ground for the theory of the Denjoy integral. Chapter 9 contains

a very careful and complete discussion of the problem of unique determination of functions of various classes by means of their derivatives or derived numbers of various sorts. Chapter 10 is devoted to the integral of Denjoy. The author gives here a remarkably clear and concise exposition of this delicate theory which he developed in his memoir published in the Fundamenta Mathematicae. Chapter 11 contains a careful exposition of some most recent results of the theory of the approximate and total differentials of a function of two variables. Some of the results which are discussed here are entirely new. The end of this chapter is of a particular significance. It contains a beautiful proof by Menchoff (never published before) of the theorem to the effect that if $f(z)$ is continuous in a domain and admits of partial derivatives finite everywhere (or even only at all points except for a denumerable set), and if these derivatives satisfy the Cauchy-Riemann equations, then $f(z)$ is analytic. This discussion represents a striking example of what can be achieved in most classical problems of analysis by methods of the modern real function theory. The last Chapter is devoted to a rapid but fairly complete exposition of the theory of the Lebesgue integral in abstract spaces. This theory should have found its way long ago into the treatises on the real function theory. The introduction of such a chapter in Saks's book is a welcome innovation. At the end of the book there is an interesting note by Banach containing a simplified exposition of the theory of measure of Haar, which proved to be of extreme importance for the theory of continuous groups. The book is supplied with a long (12 pp.) bibliography, interesting in itself. The book is very well printed and is practically free from misprints (except for those corrected in the Errata). In conclusion we may state our conviction that Saks's book is eminently suited to be used in a graduate course on the real function theory. This opinion has been checked by the reviewer's personal experience.

J. D. TAMARKIN

SHORTER NOTICES

The Expanding Universe. By Sir Arthur Eddington. New York, Macmillan, 1933. x+182 pp.

This volume, an enlarged version of a lecture delivered by Eddington at the meeting of the International Astronomical Union at Cambridge (Mass.) in Sept. 1932, treats the theory of the expanding universe, not as an end in itself, but as a means for determining the cosmical constant which appears in the Einstein theory. The subject lies at the meeting point of astronomy, relativity, and wave-mechanics. If the theory developed is valid, then (p. 170) "to measure the mass of an electron, a suitable procedure is to make astronomical observations of the distances and velocities of the spiral nebulae." This indicates the marked interest of the book for all those who delight to meditate on the interrelatedness of natural phenomena.

R. D. CARMICHAEL

Fourier'sche Reihen, mit Aufgaben. By J. Wolff. Groningen, P. Noordhoff, 1931. 60 pp.

This monograph, written in synoptic form, is an excellent condensation of material that is usually developed in many more pages. The choice of material is also commendable; the theorems selected are certainly among the most fundamental in the vast number of results concerning Fourier series available in classical and modern literature.

The book is divided into three sections, each accompanied by a valuable set of exercises. Many of the exercises, indeed, involve fundamental results in the theory of Fourier series which are sufficiently closely related to theorems in the text to justify having their development left to a student possessing initiative. The first two sections of the book can be read by those familiar only with the Riemann integral. The third section presupposes a knowledge of the theory of the Lebesgue integral.

Among results presented in the first section are criteria for convergence of the Fourier series for functions satisfying Lipschitz conditions, and functions which are monotonic, or of bounded variation, both in connection with convergence at a point and uniform convergence in an interval. There is also given an example of a continuous function whose Fourier series diverges at a point.

The second section contains a discussion of summability $(C1)$ of the Fourier series, the best approximation property, in the sense of least squares, of the Fourier partial sums, Fejér trigonometric polynomials, Parseval's theorem, some of Riemann's theory of trigonometric series in general, and some of Cantor's work on the uniqueness of Fourier developments. The Fourier integral is treated in the exercises.

The third section begins with the extension of classical results to Lebesgue integrable functions. Then, after certain properties of such functions are discussed, some of the recent results depending essentially on Lebesgue integration are obtained. There is a discussion of the convergence criteria due to de la Vallée Poussin, Lebesgue, and W. H. Young, and some of Fatou's results in

connection with the Poisson integral. The Riesz-Fischer theorem and other important results are given in the exercises.

The book as a whole should certainly serve as an excellent adjunct to a course in Fourier series, both for students of pure mathematics and of mathematical physics. It could also serve well as a guide to a student attempting to make progress in the theory by means of individual study. In this case it would be advisable to use it in connection with other texts where the treatment is less condensed. Finally, it is an excellent reference work for those already familiar with the theory who desire a compact summary of the most outstanding results.

C. N. MOORE

Einführung in die Theorie der kontinuierlichen Gruppen. By Dr. Gerhard Kowalewski. (Mathematik und ihre Anwendungen, Band 9.) Leipzig, Akademische Verlagsgesellschaft, 1931. 10+396 pp.

This attractive volume is a distinctly worth while addition to the literature of the Lie theory of continuous groups. The author, himself an erstwhile student of Lie's, has lectured at various times on this subject at a number of German universities, and from these courses, and from a basis of notes and recollections of conversations with Lie, the present work is professed to have sprung. The presentation is made with care and skill and is distinctly readable. With the introduction of each new concept the author avails himself of illustrative material from geometry or mechanics to help set forth the idea itself preparatory to its formulation in analytical garb. Thus, at the very beginning a consideration of the steady two-dimensional flow of a fluid is made to yield the concepts of the group of infinitesimal transformations, the path curves, the integrated finite transformations, etc. Critical remarks are freely interspersed in the text and add life to the presentation. At appropriate intervals the theory is summarized, and is consolidated now and again by detailed and sometimes lengthy discussions of important and interesting special cases.

Despite the smoothness of its exposition the book requires of the reader a considerable measure of mathematical maturity. A familiarity with the analysis necessary and incidental to the discussion is assumed without question. The work is not to be thought of, therefore, as one designed or suited to give a hasty and superficial acquaintance with the subject. On the contrary it seeks to present the theory in as complete a form as possible, and to this end the author has sought to incorporate with the fundamental work of Lie the important results of such modern investigators as Cartan and F. Schur.

The volume is one of 394 pages and is divided into four chapters. The first of these, entitled *Infinitesimal Transformations and One-Parameter Groups*, begins with the fundamental definitions and extends through a discussion of the integration of Lagrangian and Pfaffian systems of equations. The remaining chapters are given respectively to *Multiple-Parameter Groups and Their Infinitesimal Transformations*, to *Lie's Fundamental Theorems*, and to *Groups of Transformations on the Line and in the Plane.*

The book will be indispensable to workers in its field, and is generally to be recommended.

R. E. LANGER

Vector Analysis. By H. B. Phillips. New York, John Wiley and Sons, 1933. viii+236 pp.

The following quotation from the preface will give some idea of the contents: "In broad outline the book consists of two parts. The first five chapters cover the fundamental operations and the more general properties of scalar and vector fields. The remaining chapters contain the detailed analysis of fields, the properties of potentials, and linear vector functions. In an elementary course the work might be restricted mainly to the first five chapters together with selected topics from the others." The first part includes, in addition to chapters on the algebra and calculus of vectors, a discussion of line, surface, and volume integrals, Stokes's theorem, the divergence theorem, and generalized coordinates. The last five chapters contain applications to electricity and hydrodynamics, an introduction to potential theory, and a chapter on dyadics. Although the material covered can hardly be classified as easy for an undergraduate, the author has succeeded in making his presentation as elementary as the nature of the subject matter allows. There are 235 problems. We believe the book contains excellent material, well presented, for a semester's work following advanced calculus.

The following comment is not to be taken in any way as a criticism of the book under review. It is our opinion that the usefulness of the Gibbs vector notation is much overrated. For most purposes the representation of a vector by a typical component, using the symmetric subscript notation and the double index summation convention, is far more efficient. For an exposition of this notation the reader is referred to the recent book, *Cartesian Tensors*, by Jeffreys, or to *Vector Analysis and Relativity*, by Murnaghan.

C. A. SHOOK

Differentialgeometrie. By L. Bieberbach. Teubner's Mathematische Leitfaden, Volume 31. Berlin, Teubner, 1932. vi+140 pp.

I quote from the preface: The purpose of this "Leitfaden" is to give an introduction to the differential geometry of real curves and surfaces of the euclidean plane and euclidean space. The endeavor is made throughout to do this with the least possible prerequisite knowledge. In general only the elements of the differential and integral calculus and of analytic geometry are necessary, the last indeed in vector form . . . . "Ich hoffe weiter, dass man in meinem Buche kein saloppen Gedankengänge finden wird, keine unsauberen Schlüsse von der Art wie sie auch die moderne Literatur über Differentialgeometrie leider so oft noch beherbergt—als Überreste aus der Plüschmöbelziet und als Verstandesschoner."

The book is an extraordinarily good, rigorous, and short presentation of the principal elementary results of differential geometry with the inclusion of a good deal of interesting matter not usually to be found in a first book on the subject. There are only 132 pages of text, which are divided into three chapters: 1. *Curves in the Euclidean Plane* (pp. 1–29); 2. *Curves in Euclidean Space* (pp. 30–46); 3. *Surfaces in Euclidean Space* (pp. 47–132). The last ten pages of the third chapter give an introduction to the tensor calculus.

The presentation of the subject is clear and careful; much more attention is paid to rigor than is customary in books on Differential Geometry. The treat-

ment is extremely compact, a fact mostly due doubtless to the use of vector methods. The author has a light touch and frequently brings a smile to the face of the reader by his sarcastic comments on other (unnamed) writers. I think the book a very good one for anyone familiar with the subject, but that it should serve a beginner as a "leading string" seems to me not possible; that a student with the slight mathematical knowledge supposed to be necessary for the study of the book should be able to learn the subject from it or to acquire an interest in it does not seem likely. The discussion is somewhat uneven, often rather difficult; again, when the author remembers that he is writing a "Leitfaden," of extreme simplicity. I question whether the use of the vector method is the best way to begin the study of differential geometry, whether the use of machinery almost inevitably strange and difficult for a student of modest equipment will not distract his mind from the great geometrical interest of the subject. Without doubt the method is generally short and elegant, but there are occasions where to me it seems forced and clumsy. Moreover I am not convinced that the vector method furnishes a very good introduction to the tensor calculus; I am doubtful whether an introduction to the tensor calculus should appear in a "Leitfaden" in differential geometry, for I am not yet persuaded that the chief purpose of classical differential geometry is to serve as an introduction to the geometry of n dimensions.

Certainly there is little in the book with which one may find fault, but I do deprecate the writing of the Christoffel symbols in their historic forms when the summation convention is to be used. This convention is common and is very useful; it is introduced in the usual way on page 47 of the book. The two Christoffel symbols are introduced on pages 61 and 62,—the symbol of the second kind, it may be noted in connection with the second footnote of page 25, in a highly artificial way. The symbols are used continually with the summation convention throughout the remainder of the book. As the symbols are written the upper and lower indices are in total confusion. I think that Bieberbach should follow L. P. Eisenhart in his change in the writing of the symbols, as given in his *Riemannian Geometry* on page 17.

On pages 92–95 there is a long proof of the very interesting Gauss-Bonnet integral formula, generally omitted in elementary texts, with applications in the following pages. To this proof is appended in a footnote heavy sarcasm to the effect: "I do not know whether one may conclude that those authors have a proof which is so short that it may be left to the readers of a textbook." I hasten to state that I do not question the essential correctness of Bieberbach's proof, but he is, in his proof, chiefly concerned with the evaluation of an integral, $\int d\omega$, where ω is the angle measured from the vector tangent to a parametric curve to the vector tangent to the curve along which the integral is taken; the integral is found by the integration of $d\omega/ds$ with respect to s, the arc. The result is immediately applied to a vertex where ω changes but ds is zero. In the same proof on page 94 is the statement: "the total angle turned by the tangent as it (the point of contact) describes the curve, is, from considerations of continuity already used, equal to the total angle turned by a chord joining points sufficiently near." I can not help thinking of plush furniture. But I think too that this is a very excellent book.

J. K. WHITTEMORE

Exercices d'Analyse. By Gaston Julia. Paris, Gauthier-Villars. Vol. 2, 1932, iv+344 pp.; Vol. 3, 1933, iv+287 pp.

These two additional volumes of exercises in analysis continue in the same spirit as that of the first volume which was reviewed by E. Hille in this Bulletin, vol. 35, p. 739. Indeed the method adopted for all three books is the same, namely that of treating (solving) each of a relatively small number of problems in several very distinct ways and of comparing the different methods employed in these solutions. The volumes constitute a set of varied exercises which figure in the programme of certification of the differential and integral calculus at the École Normale Supérieure, the Sorbonne, and other schools. They in no sense replace the material of Goursat's *Cours d'Analyse*, but are, rather, companion volumes designed primarily for those students preparing for the examinations of the *licence* and the *aggrégation*. Some of the exercises are new while others have been taken from past examinations. In volume 2 (analysis), Chapters 2 and 4 and parts of Chapters 1 and 5, being descriptive in character, are more like the standard text-book. The third volume (differential equations) is given over wholly to the 45 problems contained therein.

Most of the problems have at least two solutions, some have as many as four. The results as well as the methods employed are made use of in applications which accompany the solutions; and there are such applications or remarks associated with almost every exercise. The tables of contents themselves number 12 pages and 17 pages, respectively. It would not be feasible, therefore, to consider here and in detail the variety of topics discussed; but the chapter headings of the two volumes will give some indication of the scope of the works. These are, in condensed form: vol. 2, (1) *Analytic functions*, (2) *Development in series*, (3) *Residues*, (4) *Analytic transformations*, (5) *Conformal mapping*; vol. 3, (1) *Methods of integration* (in which are treated integration by quadratures, Riccati's equation, etc.), (2) *Linear equations* (with special reference to singular points, transformations, etc.), (3) *Singular integrals*.

It is evident that the high standards of achievement required of the French student of mathematics can be matched by no such requirements even in the very best of our American universities.

C. O. Oakley

Lehrbuch der höheren Mathematik für Universitäten und Technische Hochschulen. By Gerhard Kowalewski. Berlin and Leipzig, Walter de Gruyter, 1933. Vol. I, 208 pp., Vol. II, 240 pp.

These two volumes will be followed by a third; the three volumes were prepared by Professor Kowalewski (Dresden) after nearly thirty years teaching experience. Beginning with analytic geometry he develops a wide variety of mathematical topics following an order of arrangement and methods of presentation which are not customary to American texts. In the first volume, *Vektorrechnung und Analytische Geometrie*, after defining vectors, vector operations, and vector properties, he uses vector notions to develop the theory and use of homogeneous coordinates for the points of a plane and of a line, the transformation of coordinates, principal properties of determinants, some geometric properties of triangles and quadrilaterals, homogeneous and non-

homogeneous systems of linear equations, and anharmonic ratio. After some applications to spherical trigonometry and to statics he discusses without using vectors the following topics in the order named: rotation in the plane, coaxial quaternions and ordinary complex numbers, spirals, linear fractional transformations, reflection on a circle and on a line, fundamental geometric forms of the first kind and projective relations between them, involutions on one-dimensional forms, Pascal's theorem, projectivities between planes, projective generation of conics, etc.

The second volume has two divisions. In the first, *Hauptpunkte der Analytischen Geometrie des Raumes*, the projective properties of ordinary space and of quadric surfaces are treated; in the second part, *Grundbegriffe der Differential- und Integralrechnung*, rigorous treatments are given of convergent sequences, limiting values, the operation of passing to a limit, and the derivative and differential of a function. After computing the derivatives of some elementary functions, he studies the Leibnitz fundamental theorem, namely, $dF(u)=F'(u)du$, where $u=f(x)$, then the mean value theorem, and closes with the use of integration to determine areas of plane surfaces and lengths of curves.

Since the book is designed for *Hochschulen* as well as for *Universitäten* the reviewer assumes that the student would have no previous knowledge of the subjects treated and believes that these books alone would not give an adequate knowledge of analytic geometry or of projective geometry or of the calculus, though they might be used to advantage to supplement texts which are more elementary and at the same time more complete, or to follow a first and more complete treatment of the subjects enumerated. There are few examples and no exercises for the student, nor is there a prefatory word from the author to give a clue either as to the mathematical maturity of the students for whom his books are intended or as to the length of time he would expect them to spend on this amount of material.

MAYME I. LOGSDON

Krise und Neuaufbau in den exakten Wissenschaften. Fünf Wiener Vorträge. Leipzig and Wien, Deuticke, 1933. iv+123 pp.

This volume consists of five general addresses on the present state of physics, with authors and titles (in translation) as follows:

Hermann Mark: The shock of classical physics due to experiment.
Hans Thirring: The transformation of the system of concepts of physics.
Hans Hahn: The crisis in outlook (Anschauung).
Georg Nöbeling: The fourth dimension and curved space.
Karl Menger: The new logic.

R. D. CARMICHAEL

NOTES

The founding of an international mathematical journal, Compositio Mathematica, is announced by its publisher, N. Noordhoff, of Groningen. The administrative board of editors consists of L. Bieberbach, of Berlin, L. E. J. Brouwer, of Amsterdam, T. De Donder, of Brussels, G. Julia, of Paris, and B. M. Wilson, of Dundee. They are assisted by nearly fifty editors, representing all the principal mathematical countries, including the following Americans: L. P. Eisenhart, E. Hille, S. Lefschetz, and O. Veblen.

Arrangements have been completed for the publication, through the offices of the American Institute of Physics, of the Journal of the Aeronautical Sciences, the official organ of the newly established Institute of Aeronautical Sciences. The Journal will be edited by the president of the Institute, Professor J. C. Hunsaker, head of the department of mechanical engineering at the Massachusetts Institute of Technology. The business of the Institute of Aeronautical Sciences, including the handling of membership applications and subscriptions to the Journal, will be carried on at its office, 5431 R. C. A. Building, Rockefeller Center, New York City.

An edition of the complete works of H. A. Lorentz is being prepared, under the editorship of P. Zeeman, P. Ehrenfest, and A. D. Fokker. The edition will be in nine volumes, and will include all treatises that have not already appeared in book form. The publisher is Martinus Nijhoff, of the Hague.

At a meeting held at Centenary College, Shreveport, Louisiana, on October 21, 1933, the Southern Intercollegiate Mathematics Association was founded, "to promote and support contests in mathematics and to encourage mutual and intellectual fellowship among its members." The following officers were elected: president, I. Maizlish, of Centenary College; vice-president, J. E. Burnam, of Simmons University; secretary-treasurer, Frances White, of Louisiana Polytechnic Institute.

The Nobel prize in physics for 1932 has been awarded to Werner Heisenberg, of Leipzig; the award for 1933 has been divided between Erwin Schrödinger, of Berlin, and P. A. M. Dirac, of Cambridge.

Professors F. Paschen, formerly director of the Reichsanstalt, Berlin, A. Sommerfeld, of the University of Munich, and R. W. Wood, of Johns Hopkins University, have been elected honorary fellows of the Physical Society of London.

The University of St. Andrews has conferred an honorary doctorate on Emeritus Professor J. E. A. Steggall, who has recently retired after fifty years of service in the chair of mathematics in University College, Dundee.

On the occasion of the celebration of the fiftieth anniversary of the founding of the Graduate School of the University of Pennsylvania, Dr. L. P. Eisenhart,

dean of the Graduate School of Princeton University, received the honorary degree of doctor of science.

J. LeRoux, professor of rational mechanics at the University of Rennes, has retired with the title of honorary professor.

Professor Alfred Loewy, of the department of mathematics of the University of Freiburg, has retired.

Professor L. C. Karpinski, of the University of Michigan, will lecture on various topics in the history of mathematics at the Universities of Athens, Cairo, Beirut, and Jerusalem, during the first half of the present academic year.

After attending the International Congress on Statistics held in Mexico, Professor Maurice Fréchet, of the University of Paris, delivered lectures at a number of universities in the United States. On November 2–3, he lectured at Columbia University on *The theory of dependent probabilities.*

Professor Hermann Weyl, formerly of Göttingen and recently appointed to a professorship at the Institute for Advanced Study, delivered the following series of lectures at Swarthmore College in October and November, 1933: *Subjective elements in sense perception*; *World and consciousness*; *Constructive character of scientific concepts and theories*; *Relativity*; *Subject and object in quantum physics.*

In addition to those whose names were given in a note in the September, 1933, issue of this Bulletin (p. 664), the following persons are in residence during the present academic year in the school of mathematics of the Institute for Advanced Study at Princeton: Dr. R. L. Echols, Dr. D. H. Lehmer, Dr. Mabel F. Schmeiser, Dr. I. J. Schoenberg, Dr. Anna A. Stafford.

Dr. T. C. Benton, of Pennsylvania State College, has been promoted to an assistant professorship.

Dr. Richard Brauer, formerly of the University of Koenigsberg, is spending the present year at the University of Kentucky as visiting professor of mathematics. In the November, 1933, issue of this Bulletin, p. 859, his name was given erroneously as Felix Brauer.

Dr. R. S. Burington has been promoted to an assistant professorship of mathematics at the Case School of Applied Science.

Assistant Professor H. B. Curry, of Pennsylvania State College, has been promoted to an associate professorship.

Professor James Franck, formerly of Göttingen, was the joint guest of the physics departments of Harvard University and the Massachusetts Institute of Technology during the month of December, 1933.

Associate Professor Orrin Frink, of Pennsylvania State College, has been promoted to a full professorship of mathematics.

Mr. W. H. Ingram has been appointed lecturer at the College of the City of New York.

The Abbé Georges Lemaitre, professor of astrophysics at the University of Louvain, has been appointed visiting professor at the Catholic University of America for the present academic year.

Professor John von Neumann, of Princeton University, has been appointed professor of mathematics at the Institute for Advanced Study.

Dr. H. H. Pixley, of the College of the City of Detroit, has been granted leave of absence to serve as mathematical economist in the Division of Economic Research and Planning of the National Recovery Administration.

Associate Professor C. A. Rupp, of Pennsylvania State College, has been promoted to a professorship of mathematics.

Assistant Professor I. M. Sheffer, of Pennsylvania State College, has been promoted to an associate professorship.

Associate Professor C. C. Wagner, of Pennsylvania State College, has been promoted to a professorship.

The following appointment to an instructorship in mathematics is announced:

Johns Hopkins University: Dr. J. L. Dorroh.

The widow of the late Dr. J. C. Glashan, distinguished Canadian mathematician, desires to dispose of his library, which consists of about five thousand volumes, chiefly modern treatises on pure and applied mathematics. Contact with the owner may be made through Professor James T. Shotwell, of the Carnegie Endowment for International Peace, 450 West 117th St., New York City.

Dr. P. Ehrenfest, professor of theoretical physics at the University of Leiden, committed suicide on September 25, 1933.

P. Forchheimer, formerly professor at the Technical School of Graz and later director of the Technical School at Constantinople, died at the age of eighty-two.

Mr. E. B. Hill, assistant actuary of the Aetna Life Insurance Company, died October 13, 1933, at the age of forty-eight.

Professor A. B. Morton, of the Georgia School of Technology, died October 13, 1933. Professor Morton was a member of the American Mathematical Society.

Miss Mary F. Smith, College Recorder of Wellesley College, is dead. Miss Smith was a member of the American Mathematical Society.

ABSTRACTS OF PAPERS

SUBMITTED FOR PRESENTATION TO THE SOCIETY

The following papers have been submitted to the Secretary and the Associate Secretaries of the Society for presentation at meetings of the Society. They are numbered serially throughout this volume. Cross-references to them in the reports of the meetings will give the number of this volume, the number of this issue, and the serial number of the abstract.

1. Professor D. N. Lehmer: *A law of reciprocity for the entries in a table of linear forms.*

In constructing tables of linear forms for use in making his factor stencils the author found the following theorem of prime importance: If A and B are entries in any table, so also is the product AB an entry. This theorem suggested a close analogy between the theory of the entries in these tables and the theory of quadratic residues. This analogy is far reaching. It leads to the following analog of Legendre's Law of Reciprocity: If P and Q are two odd primes, then P is an entry in the table for Q whenever Q in an entry in the table for P unless P and Q are both of the form $4n-1$, in which case if P is in the table for Q, Q will not be in the table for P. Easy modifications are necessary for negative values of P and Q. Linear form tables may also be constructed giving the tables which have a given entry. (Received November 6, 1933.)

2. Professor Morgan Ward: *The diophantine equation* $X^2-DY^2=Z^M$.

All relatively prime integral solutions of the diophantine equation $X^2-DY^2=Z^M$ are obtained by the use of ideals under the assumptions that D is square-free, incongruent to one modulo 8, while M is any positive integer prime to the class number of the quadratic field $K(D^{1/2})$. The formulas correct and extend earlier attempts by Pepin and others to solve the equation, and are applied to discuss the solution of various allied diophantine equations. (Received November 8, 1933.)

3. Dr. D. C. Duncan: *A plane rational curve of order* $2k+1$.

In an earlier paper (to appear in a forthcoming issue of this Bulletin) the writer has established the existence of real, non-degenerate, completely symmetric, self-dual, *elliptic* curves of order $4k+2$, with the singularities all distinct, together with a plan for sketching them approximately. In the present paper one observes that the completely symmetric, self-dual *rational* curve with its singular elements all distinct has a real existence for all orders. Moreover, a very good approximation of such a locus of order n (i.e., $2k+1$) is realized by drawing secant lines through all *consecutive* pairs of $n-2$ points

equally distributed around a circle, omitting the chords. The $n-2$ acnodes appear within the circle between consecutive cusps. These rational curves are also invariant under $4k-2$ collineations and $4k-2$ correlations, of which $2k$ are polarities, all of which are listed. The $(2k+1)^2$ foci are shown to admit fairly easy determination. The equations in Cartesian rectangular coordinates are given for the curves of orders 5, 7, and 9, together with a sketch of the locus of order 9, depicting all the singular elements and autopolarizing conics. (Received November 10, 1933.)

4. Professor A. R. Williams: *The apparent contour of the general V_3^n in S_4.*

It is the purpose of this paper to study the surface obtained by intersecting with a 3-space the hypercone of tangents drawn to a hypersurface V_3^n in S_4 from a point O. The two cases when O is not a point of V, and when it is on V, are considered in that order. Results obtained are checked by the known formulas for surfaces in S_3. (Received November 10, 1933.)

5. Professor R. S. Burington: *A classification of plane cubic curves under the affine group by means of arithmetic invariants.*

In this paper a system of arithmetic invariants is exhibited which is sufficient to give a complete separation of real ternary cubics, under the real affine group, into 110 species and 42 canonical forms. Associated with the cubic, C, is a certain set of matrices whose ranks and signatures serve to make the division into species. A prominent part is played by the matrix of the quadratic hessian of C, and by certain quartic covariants and their discriminant matrices. The appearance of certain families of cubics in recent electrical work should add interest to the results of this paper. (Received November 8, 1933.)

6. Mr. P. M. Pepper: *Affine-polygenic functions.*

That the real and imaginary parts of $w(z)=u(x, y)+iv(x, y)$ satisfy at every point in a given region R a pair of partial differential equations of the type $u_x+a_2v_x-a_1v_y=0$, $-v_x+a_2u_x-a_1u_y=0$, a_1, a_2 real, $a_1\neq 0$, characterizes all functions $w(z)$ analytic throughout R in some variable $Z=X+iY$, $X=a_{11}x+a_{12}y+a_{13}$, $Y=a_{21}x+a_{22}y+a_{23}$, a_{ij} real, $a_{11}a_{22}-a_{21}a_{12}\neq 0$. The particular transformation $X=a_1x+a_2y$, $Y=y$ will always yield analyticity. Given $w(z)$, substitution of the specific functions u_x, u_y, v_x, v_y in the above differential equations and solution for a_1 and a_2 in terms of these partial derivatives affords a test for transformability and if $w(z)$ is thus transformable, yields, simultaneously, the constants a_1 and a_2 of a transformation. Integrals of such functions about closed contours are expressible as line integrals and as area integrals of analytic functions: $\int w(z)dz=\pm\frac{1}{2}K\int f(Z)(dX-idY)=\mp iK\iint f'(Z)dXdY$, $f(Z)$ and $f'(Z)$ being, respectively, the transform of $w(z)$ and the derivative of this transform with respect to Z. The first two integrals are taken about, respectively, the original contour and the transformed contour, the third integral is taken over the area enclosed by the transformed contour. The constant K has the value $(1-a_1-ia_2)/a_1$. The upper or lower signs are taken according as a_1 is positive or negative. (Received November 8, 1933).

7. Professor Louis Brand: *A significant vector in the theory of surfaces.*

If a portion of a surface is simply covered by a field of curves of unit tangent vector a, we may form at every point a unique trihedral anb, where n is the unit surface normal and $b=a\times n$. As the point P moves with unit speed along a surface curve of unit tangent t, the angular velocity Ω of the trihedral anb is a function of the position of P and of the direction t. In fact $\Omega=t\cdot\Phi$, where Φ is a dyadic function of position. This dyadic may be expressed as $\Phi=Qn-\nabla n\times n$, where ∇n is the surface gradient of n. This paper deals with the properties of the vector Q thus defined. The integrability conditions for the surface are summarized in $n\cdot\operatorname{rot}\Phi=n\cdot\Phi_2$, where Φ_2 is the second of Φ in the sense of Gibbs. In particular $n\cdot\operatorname{rot}\nabla n=0$ gives the Codazzi equations, while $n\cdot\operatorname{rot}Q=-K$ gives the equation of Gauss, and at the same time is the vector expression of Gauss' *Theorema Egregium* on the total curvature K. The integral form of this equation, namely $\int K\,d\sigma=-\oint t\cdot Q\,ds$ is precisely *Bonnet's Integral Formula.* (Received November 7, 1933.)

8. Professor C. G. Latimer: *On the units in a cyclic field.*

In this paper three theorems are proved on the existence of certain types of systems of fundamental units in a cyclic field F. If F is imaginary, it is shown that there is a fundamental system in which every unit is real. If F is obtained by composition of two fields F_1 and F_2, it is shown that a fundamental system in F_1 together with a fundamental system in F_2 and certain other units form a fundamental system in F. In another paper the writer expressed the class number of F in terms of an ideal in a certain commutative ring. It is shown that F contains a unit such that certain of its conjugates form a fundamental system if and only if this ideal is a principal ideal. (Received November 10, 1933.)

9. Mr. H. E. Vaughan: *Some theorems on connected sets.*

This paper contains theorems on connected sets and sets irreducibly connected about subsets. Some of these are generalizations of theorems in sections one and two of the paper *Sur les ensembles connexes* by Knaster and Kuratowski (Fundamenta Mathematicae, vol. 2). (Received November 9, 1933.)

10. Professor W. L. Ayres: *On the relation between local connectivity and another property.*

Kuratowski and Hahn have shown that a continuum is locally connected at every point if and only if for each point p every neighborhood of p contains a connected neighborhood of p. In this note these properties are considered for a single point p of the continuum and it is shown they are not equivalent. This question arose in connection with a conjecture of W. A. Wilson (American Journal of Mathematics, vol. 54, p. 382), which is shown to be false. (Received November 9, 1933.)

11. Professor H. S. Pollard: *On the relative stability of the median and arithmetic mean.*

In this paper the relative stability of the median and arithmetic mean of frequency distributions which may be dissected into two and three normal distributions is investigated. The measure of stability employed is the standard deviation of these averages. It is shown that the formula which is ordinarily used to calculate the standard deviation of the median yields only an approximation to its actual value, and that for samples which do not contain a fairly large number of items and for distributions in which the median is located at a point of relatively small frequency this approximation may be untrustworthy. A second method of determining the relative stability of the two averages is then developed. The frequency functions according to which the arithmetic means and medians of samples drawn from a given distribution are distributed are obtained, and the deviations of corresponding percentiles of the two averages are compared. Both methods of determining the relative stability of the median and mean are applied to a particular sequence of economic data. (Received November 27, 1933.)

12. Dr. R. H. Cameron (National Research Fellow): *Algebraic functions of Stepanoff almost periodic.*

In this paper sufficient conditions are given under which a polynomial equation with Stepanoff almost periodic coefficients will have Stepanoff almost periodic solutions. (Received November 8, 1933.)

13. Professor Lincoln La Paz: *On a lemma of Fejér.*

L. Fejér (Mathematische Annalen, vol. 61 (1905), p. 432) has verified and used the following lemma: If $y''=F(x, y, y')$ is the Euler equation of the variation problem $\int_{x_1}^{x_2} f(x, y)\phi(y')dx$, then $y''=-F(x, y, y')$ is the Euler equation of the problem $\int_{x_1}^{x_2} [\phi(y')/f(x, y)]dx$. In the present paper the following generalization of Fejér's lemma is obtained: If for a non-singular problem of minimizing an integral $\int_{x_1}^{x_2} f(x, y_1, \cdots, y_n)\phi(y_1', \cdots, y_n')dx$ with integrand function of class C'' the Euler system in normal form is $y_k''=F_k(x, y_1, \cdots, y_n, y_1', \cdots, y_n')$, $k=1, 2, \cdots, n$; then for a problem of minimizing the integral $\int_{x_1}^{x_2} \Phi(f)\phi(y_1', \cdots, y_n')\,dx$, where Φ is of class C'' but is otherwise an arbitrary function of its argument, the Euler system in normal form is $y_k''=(f\cdot\Phi'/\Phi)F_k$, $k=1, 2, \cdots, n$. The generalized lemma is of service in writing down by inspection the normal Euler equations of the important class of brachistochronic integrals $T=\int_{x_1}^{x_2} [ds/v(r)]$ by reference to the corresponding equations derived in the simple special case $v(r)=r^{-1}$. A generalization of Fejér's lemma for double integrals of the type $\iint f(x, y, z)\phi(p, q)dxdy$ is also indicated. (Received November 8, 1933.)

14. Professor G. Y. Rainich: *Singularities on maps of surfaces.*

A map is obtained by introducing on a surface and on a plane curvilinear coordinates and representing points of the surface by points of the plane with the same coordinates; the coefficients of the two fundamental forms, given as

functions of coordinates, determine the surface. Do closed curves on the map surrounding a singular point of the coefficients correspond to closed curves on the surface? This question is settled by product integrals. Assuming an affirmative answer an invariant is introduced, the number of interlinkings of two surface contours corresponding to two neighboring map contours. First the situation in the case of a ruled surface formed by the normals to a closed curve is studied. Here the invariant is, in the case of a plane contour, an integral of the expression whose vanishing means that the ruled surface is developable. This result is used to express the invariant in the general case as a contour integral involving the fundamental forms; the integrand, in the case of a plane contour, may also be described as the difference of the two principal curvatures times the sine of the double angle between the direction of the contour and a principal direction. (Received November 9, 1933.)

15. Professor I. A. Barnett: *Transformation of an integro-differential equation by means of a Volterra transformation.*

The author shows that certain classes of linear integro-differential equations may be transformed into linear differential equations by means of a Volterra transformation. This leads at once to an existence proof for the solutions of such equations. In fact, an explicit expression is found for the solution of the integro-differential equation in terms of the infinitesimal transformation which generates the finite Volterra transformation. (Received November 10, 1933.)

16. Mr. H. S. Kaltenborn: *Some properties of Stieltjes integrals.*

The definition of Stieltjes integral has been modified and extended in various ways, and different types of Stieltjes integrals have been obtained. This paper is concerned with the four types discussed by H. L. Smith (Transactions of this Society, vol. 27 (1925), p. 491–515), which may be grouped in pairs as "ordinary" and "mean" integrals according to the way in which the approximating sum is formed, the two integrals in each group being distinguished by the method used in passing to the limit. It is shown that the more important elementary properties of the ordinary integrals are also valid for the mean integrals. A fundamental theorem is established for the mean integrals, by means of which a theorem corresponding to the substitution theorem for the ordinary integrals is derived. Several related properties, which are valid for both the ordinary and mean integrals, are then derived from the substitution theorem. The paper concludes with some sufficient conditions for the existence of the mean integrals; a particular result being that the integral of $f(x)$ with respect to $\phi(x)$ exists if $f(x)$ and $\phi(x)$ are any two regular functions of bounded variation. (Received November 10, 1933.)

17. Professor C. N. Moore: *On convergence factors for double series that are summable of non-integral orders.*

In many questions of analysis it is of importance to determine how the introduction of certain factors in the terms of a series affects the behavior of

the series as to convergence, summability, or other general properties. Factors that maintain convergence for convergent series or produce convergence in the case of summable series are naturally designated convergence factors. In a previous paper (Transactions of this Society, vol. 29 (1927)) the author has obtained necessary and sufficient conditions for convergence factors in multiple series that are convergent or summable by Cesàro means of integral orders. Interesting and important applications of some of these results to the theory of multiple factorial series may be found in a recent paper by C. R. Adams (Annals of Mathematics, vol. 32 (1931)). The aim of the present paper is to extend the theorems of the 1927 paper that concern double series to the case of series that are summable by means of non-integral orders. It is found that if the double differences occurring in the conditions of the previous theorems are suitably generalized to differences of non-integral orders, the generalized theorems may be stated in essentially the same form. The proofs, however, are considerably more complicated. (Received November 7, 1933.)

18. Miss Alta Odoms: *On the Cesàro mean of double Fourier series.*

This paper is an extension to functions of two variables of some theorems proved by O. Szász concerning the Cesàro mean of the rth order $(0<r\leqq 1)$ of the Fourier development of a function of one variable. Let the function $f(x, y)$ be integrable (L) and finite in a cross neighborhood of the point (x_0, y_0) under consideration, and let the Cesàro mean of the rth order $(0<r\leqq 1)$ of its Fourier development be denoted by $S_{mn}^{(r)}(x, y)/C_{mn}^{(r)}$. Further, let α and β be quantities such that $0<\alpha\leqq r\leqq 1, 0<\beta\leqq r\leqq 1$. Various results are obtained concerning the limits of the products of $S_{mn}^{(r)}(x, y)/C_{mn}^{(r)}$ by $m^{\alpha}n^{\beta}$, $m^{r}n^{\beta}/\log m$, $m^{\alpha}n^{r}/\log n$, and $m^{r}n^{r}/\log m \log n$, respectively, as m and n become infinite. (Received November 7, 1933.)

19. Dr. E. S. Akeley: *An invariantive characterization of quadratic differential forms.*

Consider two quadratic differential forms: $g_{ij}dx^{i}dx^{j}$ and $h_{ij}dx^{i}dx^{j}$ $(i, j=1, \cdots, n)$. Let $(\lambda_1, \cdots, \lambda_n)$ be the characteristic values of λ for the pencil $h_{ij}-\lambda g_{ij}$. If $\partial(\lambda_1 \cdots \lambda_n)/\partial(x_1 \cdots x_n)\neq 0$, then $(\lambda_1, \cdots, \lambda_n)$ will form a canonical coordinate system for this pair of forms. If, furthermore, h_{ij} be derived from g_{ij}, then $(\lambda_1, \cdots, \lambda_n)$ can be considered as a canonical coordinate system for $g_{ij}dx^{i}dx^{j}$. The case where h_{ij} is the contracted Riemann-Christoffel tensor for g_{ij} was considered in a previous paper (Abstract No. 39–5–147), where the properties of these forms in the neighborhood of the origin were considered. In the present paper, exact solutions corresponding to non-degenerate quadratic forms have been found. (Received November 13, 1933.)

20. Mr. E. H. Hanson: *A new proof of a theorem of Denjoy, Young, and Saks.*

It was shown by A. Denjoy (Journal de Mathématiques, (7), vol. 1 (1915), p. 105) that at every point, with the possible exception of a set of Lebesgue measure zero, the Dini derivatives of a continuous function satisfy one of the

following relations: (1) all four derivatives are finite and equal; (2) the two upper derivatives are $+\infty$, and the two lower derivatives are $-\infty$; (3) the upper derivative on one side is $+\infty$, the lower derivative on the other side is $-\infty$, and the other two derivatives are finite and equal. This theorem was shown to hold for measurable functions by G. C. Young (Proceedings of the London Mathematical Society, (2), vol. 15 (1916), p. 360). It was partially extended to unrestricted functions by S. Banach (Comptes Rendus, vol. 173 (1921), p. 457) and was first completely extended by S. Saks (Fundamenta Mathematicae, vol. 5 (1924), p. 98) who based his proof on the corresponding theorem for monotone functions (H. Lebesgue, *Leçons sur l'Intégration* (1904), p. 128). Another proof, as yet unpublished, is due to H. Blumberg who makes use of the result for measurable functions. The present proof, assuming no knowledge of the result for special cases, proceeds along the most natural and direct lines and shows that no simplification is gained by formulating the proof for a restricted class of functions rather than for the general case. (Received November 13, 1933.)

21. Professor O. E. Brown: *The equivalence of n-tuples of matrices.* Preliminary report.

In the non-singular case, the equivalence, in the field of their elements, of two n-tuples of square matrices of order m depends upon the solution of a system of $m^2(n-1)$ linear homogeneous equations in m^2 variables. The variables may be arranged as the elements of a square matrix of order m and a given solution of the system called a solution matrix. The given n-tuples are equivalent if, and only if, the system of equations has a non-singular solution matrix. (Received November 21, 1933.)

22. Dr. W. E. Maier: *Quadratic integral equations.*

F. Bernstein and S. Doetsch in 1922 established an integral addition theorem for elliptic modular functions. Generalizing those results to elliptic theta functions $\sum_{k=-\infty}^{+\infty} e^{2\pi i k z + \pi i k^2 \omega} = \theta(z, \omega)$ it was found convenient to introduce the *normalized* quantity $t^{-1/2}\theta(z, i/t) = \tau(z, t)$. The latter admits of an identical relation with respect to z and t, viz. for $0 < \Re(t)$, $\int_0^1 d\rho\, \tau(z, t-\rho t)[\tau(z, \rho t) - \frac{1}{2}\tau(0, \rho t/4) + \frac{1}{2}] = \tau(z, t)$. (Received November 20, 1933.)

23. Mr. W. H. Ingram: *On the dynamical theory of electrical commutator machines.*

Following Kron, we use tensor notation and write the equations of motion for a slip-ring machine in the form (1): $e_i = R_{ij}\dot{q}^j + L_{ij}(d/dt)\dot{q}^j + \Gamma_{i;mn}\dot{q}^m\dot{q}^n$, where $\Gamma_{i;mn}$ is the ordinary Christoffel symbol and where the coordinates are true. A transformation to quasi-coordinates, (2): $\dot{\xi}_i = \alpha_j{}^i(\theta)\dot{q}^j$ $(j=1, 2, \cdots, n-1)$, $\dot{q}^j = \beta_k{}^j(\theta)\dot{\xi}^k$ $(k=1, 2, \cdots, n-1)$, $\xi_n = q^n = \theta$, where θ is the privileged rotor position coordinate, leads to the system (3): $e_\sigma = r_{\alpha\sigma}\dot{\xi}^\alpha + l_{\alpha\sigma}(d/dt)\dot{\xi}^\alpha + \lambda_{\sigma;\beta\gamma}\dot{\xi}^\beta\dot{\xi}^\gamma + l_{\beta\alpha}\beta_{k\alpha}\beta_{n\gamma}(\partial\alpha_{k\alpha}/\partial\theta)\dot{\xi}^\beta\dot{\xi}^\gamma$, where $\lambda_{\sigma;\beta\gamma}$ is also a Christoffel symbol of the first kind. The system (3) is known to be identical with that for a commutator machine with an infinite number of commutator segments. It is now found to be identical with the equations of motion of a Lagrangean system

whose kinetic energy is expressed in terms of the ξ's and θ and when the ξ's are regarded as moving coordinates. This suggests that the relative angular velocity of commutator and brush is a dynamical homologue of the angular velocity of a system of reference axes and that the last term in (3) may be called the Coriolis voltage. (Received November 20, 1933.)

24. Dr. G. G. Estes: *The lift and the moment of an arbitrary aerofoil Joukovsky potential.*

In this paper is taken up the problem of finding the lift and the moment of an arbitrary aerofoil in a flow which is according to Joukovsky's theory. Given an arbitrary aerofoil, we first select a Joukovsky profile which is near the given profile. We then calculate the Green's function (pole at infinity) for the arbitrary profile by use of the following formula due to Hadamard: $\delta G_\infty^A = -(1/(2\pi))\int (dG_\infty^M/dN)\ (dG_M^A/dN)\delta N ds$. G_∞^A is the value at the point A of the Green's function with pole at infinity. The path of integration is the Joukovsky aerofoil. δN is the normal variation from the Joukovsky aerofoil to the arbitrary aerofoil. A new labor-saving method of evaluating the above integral is given, and is carried through for an arbitrary aerofoil. We get the value of the new Green's function (G') at the points A by adding the variations given by the integral above to G_∞. Using a Fourier series expansion of G', we get the function conjugate to G' (call it H'). The function that maps the region exterior to the arbitrary aerofoil on the interior of the unit circle is now known, and it is now possible to get the flow about the arbitrary aerofoil. The lift and the moment are then calculated by well known formulas of aerofoil theory. (Received November 29, 1933.)

25. Mr. W. L. Morris: *A new method for the evaluation of double integrals.*

The evaluation of $\iint_A f(x, y)\, dy\, dx$ is not always possible by simple integration. In order to take care of the cases where simple integration is impractical, several methods have been devised. One is that of transformation, where a substitution is made on x and y in an attempt to simplify the limits of integration. This ordinarily renders the integrand more complicated. If, however, we place (1) $f(x, y) = J$ where J is the Jacobian of u, v with respect to x, y, it is possible to find $u = u(x, y)$ and $v = v(x, y)$ satisfying (1). Then $\iint_A f(x, y)\, dy\, dx = \iint_{A'} du\, dv$. This is a simple area integral over A', the evaluation of which may be accomplished readily. One method that suggests itself is the graphical method. Suitable transformations for some of the more common engineering integrands have been obtained. It has been found that the application of some transformations require Riemann surfaces. The evolution of suitable mapping rules for each transformation, however, will permit the use of this method without knowledge of such surfaces. (Received November 25, 1933.)

26. Dr. Wladimir Seidel: *An extension of Cauchy's theorem.*

Let C be a closed rectifiable curve (not necessarily a Jordan curve) which forms the boundary of a simply connected region R in the z-plane. Denote by $w = g(z)$ an analytic function in the region R which maps this region con-

formally on the circle $|w|<1$. The inverse function $z=h(w)$ of $w=g(z)$ is continuous in the closed circle $|w|\leqq 1$. If the circumference $|w|=1$ is described in the counter clockwise sense, the corresponding point z will describe C in a certain orientation which we call the *simple orientation* of C. Under these conditions the extension of Cauchy's theorem may be stated as follows: *Let $f(z)$ be a function analytic interior to R, and bounded there: $|f(z)|<M$, M being a positive constant. Then $f(z)$ has limiting values along almost all normals to the curve C, defining on C a bounded measurable function $f^*(\zeta)$, where ζ is a point of C, and $\int_C f^*(\zeta)\,d\zeta=0$, the integration being taken along C in simple orientation.* In the case that C is a simple Jordan curve the theorem was proved by V. V. Golubev, Master's Thesis, Moscow, 1916. (Received November 29, 1933.)

27. Dr. R. H. Cameron (National Research Fellow): *Linear almost periodic transformations.*

Almost periodic, mono-basal, and elementary almost periodic transformations are the analogues in transformations of almost periodic, limit periodic, and actually periodic functions. In a paper to appear in the April number of the Transactions of this Society, the author has shown that an almost periodic transformation can be expressed as an infinite product of permutable mono-basal transformations. In the present paper it is shown that a linear almost periodic transformation of a Banach space into itself can be expressed as an infinite product of permutable elementary almost periodic transformations. (Received November 29, 1933.)

28. Mr. Jacques Hammond: *On capacity in potential theory.* Preliminary report.

In this paper it is proved that Kellogg's definition of capacity is equivalent to the transfinite diameter definition, which in turn is known to be equivalent to several others. The corresponding two-dimensional situation is also considered, and a new definition is proposed analogous to Kellogg's three-dimensional definition. The capacities of certain sets bounded by circles are obtained. (Received November 27, 1933.)

29. Dr. Hassler Whitney: *A numerical equivalent of the four-color problem.*

Let $(p_1, q_1), \cdots, (p_m, q_m)$ be pairs of positive integers, with each $q_i>p_i$. We say they form an *admissible set* if there are no i and j such that $p_i<p_j\leqq q_i<q_j$. The following theorem is equivalent to the four-color theorem. *If $(p_1^{(k)}, q_1^{(k)}), \cdots, (p_{m_k}^{(k)}, q_{m_k}^{(k)})$ $(k=1, 2)$ are two admissible sets, and n is the largest of the numbers $p_i^{(k)}$, $q_i^{(k)}$, then there are numbers $a_1, \cdots, a_n$, each equal to 1, 2, or 3, such that $a_{p_i^{(k)}}+a_{p_i^{(k)}+1}+\cdots+a_{q_i^{(k)}}\not\equiv 0 \pmod 4$, $(i=1, \cdots, m_k;$ $k=1, 2)$.* (Received November 29, 1933.)

30. Professor R. L. Jeffery: *Derived numbers and approximate derivatives of non-measurable functions.*

The problem of the present paper has been considered for measurable functions by J. C. Burkill and U. S. Haslam-Jones, Proceedings of the London

Mathematical Society, (2), vol. 32, p. 346. In that paper they introduce the concept of λ-derivates. In a later paper, Quarterly Journal of Mathematics, vol. 4 (1933), they use this concept, together with the idea of measurability of one set with respect to another, in a study of derivatives for non-measurable functions. The present paper obtains in a simple way the possible distribution of the values of the derived numbers and approximate derivatives of arbitrary finite functions over arbitrary sets. The discussion in no way features λ-derivatives, or measurability of sets or functions, but is based on derived numbers over sets introduced by Saks, Fundamenta Mathematicae, vol. 5, pp. 98–104, and on outer measure and outer density. (Received November 25, 1933.)

31. Dr. F. W. Perkins: *The Dirichlet problem for domains with multiple boundary points.*

This paper contains a treatment of the Dirichlet problem by a method suggested to the author by Professor O. D. Kellogg shortly before his death. A concept is introduced which is somewhat analogous to the concept of "prime ends" as developed by Carathéodory (*Über die Begrenzung einfach zusammenhängender Gebiete*, Mathematische Annalen, vol. 73 (1913), pp. 322 ff.). In this way a part of the present theory relating to the Dirichlet problem may be extended, for certain types of domains, so as to render possible (in assigning boundary conditions) a discrimination between various modes of approach to a multiple boundary point. (Received November 29, 1933.)

32. Dr. Mabel Schmeiser: *Some properties of arbitrary functions.*

Let $f(x, y)$ be an arbitrary real function, P a point of the xy plane, I_{Pd} the interval determined by the lower and upper bounds of the values of $f(x, y)$ as P is approached along the direction d, and s a given straight line in the xy plane. It has been proved by Henry Blumberg that at every point P of s except possibly at the points of a denumerable set, $I_{P\alpha}$ overlaps, laps, or abuts $I_{P\beta}$ for two fixed directions α and β on the same side of s (Fundamenta Mathematicae, vol. 16 (1930), p. 77). In this paper the two directions are freed of their fixed positions and the result is proved valid for all pairs of directions on the same side of s. An example shows that for any denumerable set D on a straight line there exists a function for which D is the exceptional set. The extension of this theorem to functions of three variables, substituting a planar direction for one of the linear directions and a given plane for the given line s, gives as the exceptional set an exhaustible plane set. (Received November 29, 1933.)

33. Dr. Saunders MacLane: *Abbreviated proofs in logic calculus.*

Any system of mathematical logic must be based upon certain processes of proof which enable new true expressions to be deduced from known formulas. The simplest such processes are syllogism and inference, while substitution with an equation or with an equivalence, the construction of a normal or half-normal form for any expression, and the constructive proof of an existence theorem are other logical processes in common use in mathematics. In this paper, all these processes and many others are defined precisely, and with com-

plete generality, on the basis of an abstract analysis of the character of symbolical expressions. In virtue of these definitions the conclusion of any process is uniquely determined by its premises. Hence in giving any proof it is possible to omit all intermediate conclusions, if only the processes involved are specified. This method of abbreviating proofs makes possible a considerable condensation and clarification of the usual detailed proofs of formal logic, and also serves as an introduction to the study of plans of proofs; that is, of the principles in a proof which determine the choice of the individual steps of that proof. Such a study opens new possibilities for the application of logic to mathematics. (Received November 25, 1933.)

34. Dr. S. S. Wilks: *The independence of estimates of variance in samples from Latin square lay-outs.*

Let X_{uv} $(u, v=1, 2, \cdots, r)$ be the element in the uth row and vth column of a matrix M of observations from a normal population having variance σ^2. Let $\overline{X}_{u.}$ and $\overline{X}_{.v}$ be the means of the uth row and vth column of M respectively, and $\overline{X}$ the mean of all observations. Furthermore, let $X_{uv_{tu}}$ $(u, t=1, 2, \cdots, r)$ be any classification of the r^2 elements $\{X_{uv}\}$ such that for each value of t no two X's lie in the same row or column of M, and let $r\overline{X}_t=\sum_{u=1}^{r} X_{uv_{tu}}$. Then, if v_r, v_c, v_t, and v_e denote R. A. Fisher's estimates of σ^2 from the sets $\{\overline{X}_{u.}\}$, $\{\overline{X}_{.v}\}$, $\{\overline{X}_t\}$ and $\{X_{uv}-\overline{X}_{u.}-\overline{X}_{.v}-\overline{X}_t+3\overline{X}\}$, respectively, it is shown by means of characteristic functions that these four estimates are distributed independently of each other. Certain generalized forms of Latin square lay-outs are also considered, including the "equalized random blocks" lay-out. (Received November 25, 1933.)

35. Dr. H. L. Dorwart: *Concerning certain reducible polynomials.*

This paper discusses the existence of the reducible polynomials mentioned in Theorem 16 of the paper *Criteria for the irreducibility of polynomials*, by H. L. Dorwart and Oystein Ore (Annals of Mathematics, (2), vol. 34 (1933), pp. 81–94). The problem of finding these polynomials is shown to be equivalent to a problem that occurs in the investigation of rapidly convergent series convenient for the computation of logarithms, which in turn is known to be equivalent to a problem in diophantine analysis. (Received November 28, 1933.)

36. Professor I. M. Sheffer: *Concerning some methods of "best approximation".*

Let $\{\phi_n(x)\}$ be a sequence of functions. A "polynomial" of order n is a linear combination of $\phi_0, \cdots, \phi_n$. Let there be given a method whereby for a given function $f(x)$, to each n there is associated an essentially unique "polynomial" $T_n(x)$ of order n. There may be properties of this method which suggest, for $T_n(x)$, the use of the description "best approximation of all polynomials of order n." Interesting examples are the well known least-square method, and the "best approximation" in the sense of Widder's generalization of Taylor's series. These two examples enjoy the following property: if we inquire

into the limit of T_n, and write $T_n=T_0+(T_1-T_0)+\cdots+(T_n-T_{n-1})$, then $T_i-T_{i-1}=c_i\psi_i$, where the set $\{\psi_i\}$ is *independent of* $f(x)$, the c_i's being determined by $f(x)$. Then, $T_n=c_0\psi_0+\cdots+c_n\psi_n$, and when n is increased, terms already present are unaltered. We shall refer to this as the property of *permanence*. This paper considers a large class of methods of "best approximation" (including the two examples mentioned) having the property of permanence. We examine the question of convergence in some cases, in particular lightening some of the conditions of Widder. (Received November 27, 1933.)

37. Professor W. H. McEwen: *Polynomials of best approximation associated with certain problems in two dimensions.*

In this paper two problems are considered, both of which are concerned with the approximate representation of a function $u(x, y)$, by means of polynomials $P_{mn}(x, y)$, throughout a closed region J of the xy-plane bounded by an algebraic curve C. In the first, it is supposed that u vanishes identically on C, and the polynomials P_{mn} are defined so as to vanish identically on C and at the same time minimize the double integral over J of the rth power ($r>0$) of $|\nabla^2(u-P_{mn})|$. In the second, u is permitted to assume arbitrary values on C, and the polynomials P_{mn} are defined so as to minimize an expression involving not only the integral mentioned above but also the maximum value of $|u-P_{mn}|$ on the curve C. In both cases results are developed concerning the convergence of the polynomials P_{mn} towards the value u as m and n both become infinite. (Received November 25, 1933.)

38. Dr. D. C. Lewis, Jr. (National Research Fellow): *On the periodic motions of dynamical systems with n degrees of freedom.*

This paper is in close relation with a paper by Birkhoff and Lewis (Annali di Matematica, (4), vol. 12), in which was proved the existence of infinitely many periodic motions in the neighborhood of a periodic motion of general stable type. In both papers the critical points of a certain function defined on an n-dimensional torus and a certain symmetric square matrix (c_{ij}) of invariants of the Hamiltonian equations play important roles. In the present paper, a closer study of these yields (under suitable conditions of generality) information concerning the characteristic exponents of the periodic motions, the mere existence of which was established in the Birkhoff-Lewis paper. The results are especially precise in the case when $\sum c_{ij}u_iu_j$ is a definite quadratic form. In this case it is proved among other things that there are infinitely many periodic motions of stable type (that is, with pure imaginary characteristic exponents) near the given periodic motion. (Received November 18, 1933.)

39. Professor C. O. Oakley: *Semi-linear equations.* Part III: *Convex polygons.*

It is found that any convex polygon, treated as a closed curve, can be represented by a semi-linear equation, that is, an equation of the form $u_0+\sum_{i=1}^{n} m_i|u_i|=0$, where the m's are constants, and where $u_i=a_ix+b_iy+c_i$, $i=0, 1, \cdots, n$. For a p-gon ($p>3$), a value of $n=p-2$ is sufficient (semi-linear representations of polygons need not be unique). However, for the case

of the triangle, $p=3$, the least permissible value of n is $n=6$. (See abstracts Nos. 37-5-179 and 38-1-21, for Parts I, II, *Regular polygons* and *Irregular polygons* respectively.) (Received December 1, 1933.)

40. Professor L. W. Cohen: *Lagrange multipliers for functions of infinitely many variables*

The existence of the Lagrange multipliers at a maximum of a function of infinitely many variables subject to an infinite set of auxiliary conditions is established. Lemmas on normal infinite determinants and associated linear systems of equations are proved, one of which incidentally removes a redundancy in an implicit function theorem for functions of infinitely many variables. (Received December 1, 1933.)

41. Professor C. N. Moore: *On criteria for Fourier constants of L-integrable functions of two variables.*

In a note published in the Proceedings of the National Academy of Sciences for September, 1933, the author has given a criterion for Fourier constants of L-integrable functions which is more general than several previously known criteria. In the present paper an analogous criterion for the Fourier cos-cos coefficients of L-integrable functions of two variables is obtained. For a doubly infinite set of constants, a_{mn}, approaching zero when m and n become infinite simultaneously, or when either index becomes infinite while the other remains fixed, one further condition is sufficient that the series $\sum a_{mn} \cos mx \cos ny$ shall converge in the region $(0<x\leqq 2\pi,\ 0<y\leqq 2\pi)$ and define there an L-integrable function of x and y. If we represent by $\Delta_{r+1,r+1}a_{mn}$ the double difference of order $(r+1)$ of the given constants which begins with the term a_{mn}, our further condition is the requirement that the double series whose general term is $m^r n^r \left|\Delta_{r+1,r+1}a_{mn}\right|$ shall converge for any $r>0$. The criterion thus obtained is more general than the criteria given by the author in the second volume of the Proceedings of the International Mathematical Congress at Zurich. (Received November 29, 1933.)

42. Professors Salomon Bochner and H. F. Bohnenblust: *Analytic functions with almost periodic coefficients.*

The authors discuss the analytic functions defined by power series of the form $\sum_0^\infty f(n)z^n$, where $f(t)$ is a uniform almost periodic function of t. When $f(t)$ reduces to an exponential polynomial, $f(t)=\sum_1^N a_\rho e^{-i\lambda_\rho t}$, $G(z)$ is the rational function $\sum_1^N a_\rho(1-ze^{-i\lambda_\rho})^{-1}$, whose poles are simple and lie on the unit circle. In the general case the authors obtain formally an infinite series, which is not necessarily convergent. They show, however, that the analytic character of $G(z)$ is precisely the same as if the series were convergent. The method used consists in replacing the multiplier $(1-ze^{-i\lambda_\rho})^{-1}$ by a more suitable function, different from it only in those intervals for λ which contain no exponent of $f(t)$. In the same manner is treated the Dirichlet series $\sum f(\log n)\, n^{-s}$ and the Laplace integral $\int_0^\infty e^{-st}f(t)dt$. The results are similar in all cases; in the last, for example, the following is obtained. The analytic function $G(s)$, the continuation of $\int_0^\infty e^{-st}f(t)dt$, $\sigma>0$, is uniform. Its singular points fill out exactly the

closure of the set $\{\lambda_\rho\}$, where $-\lambda_\rho$ are the exponents of the u.a.p. function $f(t)$. Isolated singularities are simple poles and correspond to the isolated exponents of $f(t)$. (Received December 1, 1933.)

43. Professor C. R. Adams and Mr. J. A. Clarkson: *Properties of functions $f(x, y)$ of bounded variation*

We study mainly the properties of additivity, continuity, differentiability, measurability, integrability, etc., of functions $f(x, y)$ of bounded variation in the senses of Vitali, Hardy, Arzelà, Pierpont, Fréchet, and Tonelli. Let the classes of functions satisfying the respective definitions be denoted by V, H, A, P, F, and T; our results then include the following. Functions in V, H, A, P, or F are additive; if both f_1 and f_2 are in H, A, or P, $f_1 \cdot f_2$ is in the same class; if f is in H, A, or P and is bounded away from zero, $1/f$ is in the same class; these properties are not enjoyed by the unspecified classes. If f is in P (or A or H), it is continuous almost everywhere and hence R-integrable; V, F, and T contain functions everywhere discontinuous with an arbitrarily large saltus. V, F, and T contain bounded functions that are non-measurable and hence not L-integrable. If f is in P, each of its two *total variation functions* is dominated by a summable function. If only functions continuous in x and in y are admitted, the classes are related in the same way as when only functions continuous in (x, y) are admitted; if only functions upper (or lower) semi-continuous in x and in y are admitted, the classes are related in substantially the same way as when all functions are admitted (see Clarkson and Adams, Transactions of this Society, vol. 35 (1933), pp. 824–854). (Received November 12, 1933.)

44. Professor C. R. Adams and Mr. J. A. Clarkson: *On convergence in variation.*

Let $\{f_n(x)\}$ be a sequence of functions converging on $a \leqq x \leqq b$ to a limit function $f(x)$ of bounded variation; let the total variation of $f_n (n = 1, 2, 3, \cdots)$ be denoted by $T(f_n)$ and that of f by $T(f)$; then f_n will be said to *converge in variation* to f on (a, b) when and only when we have $T(f_n)$ on $(a, b) \to T(f)$ on (a, b). The results obtained in this paper include the following. Convergence in variation on (a, b), when f is continuous, implies both $f_n \to f$ uniformly on (a, b) and $T(f_n)$ on $(x', x'') \to T(f)$ on (x', x'') uniformly with respect to x', x'' for $a \leqq x' < x'' \leqq b$; the hypothesis of continuity here imposed upon f can neither be deleted nor weakened to semi-continuity. If $\{f_n\}$ converges in variation to f on (a, b) and f is $> \alpha > 0$ (or $< -\alpha < 0$), $\{1/f_n\}$ converges in variation to $1/f$ on (a, b). A set of conditions sufficient to insure convergence in variation is given, from which it follows that the partial sum of a power series converges in variation to the sum of the series in any interval interior to the interval of convergence. (Received November 12, 1933.)

45. Professor Philip Franklin: *Derivatives of higher order as single limits.*

The nth derivative of a function at a point may be determined by a single limit involving nth differences or the values at $n+1$ evenly spaced points.

However, this limit may exist without the corresponding derivative. It is here shown that if the analogous expression for unevenly spaced points be used, the existence of the limit is a necessary and sufficient condition for the existence of a continuous nth derivative. Applications to finite Taylor developments are considered. (Received November 28, 1933.)

46. Dr. E. K. Haviland: *On H-functions and the distribution functions associated with them.*

Let $x_1(t), \cdots, x_m(t)$ be real functions bounded in $(-\infty, \infty)$ and measurable in every finite interval. It may be shown that if the curve $X(t)$: $x_1 = x_1(t), \cdots, x_m = x_m(t)$ is an H-function in the sense of Wintner, that is $\lim_{T\to\infty}\int_{-T}^{T}[x_1(t)]^{n_1}\cdots[x_m(t)]^{n_m}dt$ exists for all non-negative integers $n_1, \cdots, n_m$, then there exists a unique function $\phi(E)$ representing the asymptotic relative frequency of the curve in the set E. Conversely, if $X(t)$ possesses such a distribution function $\phi(E)$, $X(t)$ is an H-function, so that the H-condition is a necessary as well as a sufficient condition that the curve possess a distribution function. This, together with the result of Birkhoff's general ergodic theorem, viz., that for dynamical systems possessing an integral invariant the H-condition is necessarily fulfilled for all solutions up to those of a space probability zero, shows that the H-condition is not as special as it appears. If $\phi(E)\neq 0$ for all sets E containing points of the curve in the m-space, $X(t)$ is called a proper H-function. Not all H-functions are proper, but if the $x_i(t)$ are almost-periodic, then $X(t)$ is a proper H-function, and this is true also of the functions considered by Birkhoff. (Received November 25, 1933.)

47. Professor V. C. Poor: *Polygenic functional solutions of certain types of integral equations.*

In this paper the uniqueness and the existence of the solution of integral equations of the Fredholm type are given. The region of integration is a complex domain or area, or a combination of both, and the solution is a polygenic function, in general non-analytic. (Received November 22, 1933.)

48. Dr. W. R. Thompson: *On the tetradite Ψ-function with application to apportionment theory.*

Defining the Ψ-function by $\Psi(r, s, r', s') \equiv [(r+s+1)!(r'+s'+1)!/(r!s!r'!s'!)]\int_0^1 x^r(1-x)^s\int_x^1 y^{r'}(1-y)^{s'}dydx$, we here show that $\binom{r'+s+r'+s'+2}{r+s+1}\cdot\Psi(r, s, r', s') \equiv \sum_{\alpha=0}^{r'}\binom{r+r'-\alpha}{r}\binom{s+s'+1+\alpha}{s} \equiv \sum_{\alpha=0}^{\alpha\leq s,\, r'}\binom{r+r'+1}{r'-\alpha}\binom{s+s'+1}{s-\alpha}$, and that Ψ is identical with the probability that drawing at random without replacements from $r+s+1$ *white* and $r'+s'+1$ *black* balls we shall draw $r+1$ *white* before $r'+1$ *black*. In an article *On the likelihood that one unknown probability exceed another in view of the evidence of two samples* (Biometrika, in press), the author has introduced a method of apportionment based on the Ψ-function. The ball-draft theorem permits the use of a machine in such apportionment which may be generalized for the case of an arbitrary number of rival *treatments*. It is further shown that $\Psi(r, s, r', s') \equiv \Psi(r, r', s, s') \equiv \Psi(s', r', s, r) \equiv 1-\Psi(r', s', r, s)$. (Received November 14, 1933.)

49. Dr. W. R. Thompson: *Recursion formulas for the tetradite Ψ and the incomplete I-function.*

The object of this paper is to establish first the relation $\Psi(r, s, r', s') \equiv \Psi(r-1, s, r', s') - (s+s'+1)\binom{r+r'}{r}\binom{r+r'}{s}/[(r+s+1)\binom{r+s+r'+s'+2}{r+s+1}]$, and secondly the relations $I_p(r+1, s+1) \equiv I_p(r, s+1) - \binom{r+s}{s} p^r(1-p)^{s+1} \equiv I_p(r+1, s) + \binom{r+s}{r} p^{r+1}(1-p)^s$, where $I_p(u, v)$ is the function of Pearson and Müller. The difference formulas for the other variables in Ψ are readily obtained by use of the permutations previously given, and the I-function formulas extended throughout their analytic continuation if the binomial coefficient symbol is generalized to include the meaning of the corresponding Gamma function ratio. (Received November 14, 1933.)

50. Dr. W. T. Reid: *An auxiliary theorem associated with the calculus of variations.*

For the problems of Lagrange, Bolza, and Mayer in the calculus of variations, the following theorem is important: If $E_{12}: y_i = y_i(x)$ $(i=1, \cdots, n;\ x_1 \leqq x \leqq x_2)$ is a normal extremal normal on every sub-interval $x_1x_2 (x_1 < x_2 \leqq x_2)$, and which satisfies the strengthened Clebsch and Mayer conditions, then there exist n mutually conjugate solutions $\eta_{ij}(x)$, $\zeta_{ij}(x)$ of the accessory equations with $|\eta_{ij}| \neq 0$ on x_1x_2. The proof of this theorem originally given by Bliss required E_{12} to have an extension normal on every sub-interval (American Journal of Mathematics, vol. 52, p. 736). Morse gave a proof for the problem of Lagrange that did not require this additional hypothesis (Annals of Mathematics, vol. 32, p. 567). Bliss and Hestenes (Transactions of this Society, vol. 35, p. 319) have used for the Mayer problem essentially the method given by Morse. In the present note this theorem is proved in a very simple manner, depending upon the method originally given by Bliss, together with a suitable modification of the accessory system in a neighborhood of $x = x_1$. The conclusion of the theorem is obtained under the weaker hypothesis that there is an integer $q \geqq 0$ such that on every interval x_1x_2 $(x_1 < x_2 \leqq x_2)$ there are exactly q linearly independent solutions η_i, ζ_i of the accessory system for which $\eta_i = 0$. (Received November 29, 1933.)

51. Professor Wilhelm Maier: *Integral equations and elliptic functions.*

F. Bernstein and G. Doetsch in 1921 found an interesting connection between modular functions and quadratic integral equations. Generalizing those results and representing the elliptic theta functions in normalized form, $t^{-1/2}\sum_\lambda \exp(2\pi izh - \pi h^2 t^{-1}) = \tau(z, t)$ for $R(t) > 0$, we find $\int_0^1 d\rho\, \tau(z, t-\rho t)[\tau(z, \rho t) - \frac{1}{2}\tau(0, \rho t/4) + \frac{1}{2}] = \tau(z, t)$. Dealing with analytic functions, defined by integral equations, the role of singular lines appears in a new aspect. Though natural boundaries for analytic continuation, they may admit the existence of certain integral mean values of the function along and even beyond the classical domain of existence. (Received November 28, 1933.)

52. Professor V. C. Poor: *Certain fundamental notions and theorems in polygenic function theory.*

This paper treats two fundamental ideas, regularity and residual values. Sufficient conditions for the validity of the circulation theorems are obtained, and an extension of the circulation theorems to an annular domain. This extension or lemma is used in proving that in the definition for the residue of a polygenic function an arbitrary contour may be used, which in the limit contracts to a point. Finally two theorems are obtained: The residue of a polygenic function, regular in a domain, D, is zero at every point of the domain. Regularity in the Stolz sense is a sufficient condition for an areal derivative in the Pompeiu sense. (Received November 22, 1933.)

53. Professor Dunham Jackson: *Note on relations connecting certain cases of convergence in the mean.*

When a sequence of approximating polynomials or trigonometric sums is determined by a criterion of closest approximation which requires the minimizing of an integral containing a power of the error, theorems on the uniform convergence of the approximation can be regarded in some cases as setting up a relationship between the convergence properties of two measures of the discrepancy between the approximating sum and the function approximated, namely the mean value of the power of the error which enters into the integral on the one hand and the maximum error on the other. This note points out that the proof of such a theorem yields at the same time a comparison of convergence properties in which the two measures of discrepancy involved are the means of two different powers of the error. (Received November 13, 1933.)

54. Professor Marston Morse: *Instability and transitivity.*

A dynamical system is said to be regionally transitive if there exists a motion whose closure is the whole of phase space. The work of Birkhoff, Hopf, Koopman, von Neumann, P. Smith, and others has thrown much light on questions of transitivity. However, it is not yet known in the analytic case whether regional transitivity implies metric transitivity. The previous examples of regional or metric transitivity have been special in character. The present paper shows that a general type of geodesic motion is regionally transitive. The geodesics considered lie on any closed Riemannian 2-dimensional manifold R of genus $p>1$. *A sufficient condition that such a system be regionally transitive is that the geodesics on R be uniformly unstable.* Instability is defined in terms of the equation of normal variation from a given geodesic. In the case where the geodesics on R are uniformly unstable the geodesics whose limit motions include all motions have the power of the continuum. The present paper defines *relative transitivity* and shows that it always exists among the minimizing geodesics of a suitable class Ω on R. When the system is uniformly unstable, Ω becomes the set of all geodesics on R, and relative transitivity becomes regional transitivity. (Received November 29, 1933.)

55. Professor G. C. Evans: *A necessary and sufficient condition for a regular point.*

Let $f(e)$ be a distribution of positive mass, finite in total amount, on a domain T, of boundary t, whose Newtonian potential is $U(M)$. The masses obtained by the sweeping out process, applied to T, according to de la Vallée Poussin converge in the weak sense to a mass $\mu(e)$, and the potential converges to a limiting function $V_0(M)$. Let the potential of $\mu(e)$ be $V(M)$. Then $V(M) \leqq V_0(M)$, but the average of $V(M)$ on any spherical surface is the same as that of $V_0(M)$. Hence it may be deduced that a necessary and sufficient condition that a point Q of t be regular for T is that $V(Q) = V_0(Q) = U(Q)$ for arbitrary $f(e)$. A sufficient condition is a similar equation, where $f(e)$ is a single point mass. (Received December 1, 1933.)

56. Professors Einar Hille and J. D. Tamarkin: *On the theory of Laplace integrals.* II.

Starting from the theory of multiplication of Laplace integrals the authors derive a simple method of analytic continuation of such functions based upon representations of the form $[\mu(z)]^{-1}\int_0^\infty e^{-zu}\,dA_\mu(u)$. Here $\mu(z) \neq 0$ is an absolutely convergent Laplace integral. The choice $\mu(z) = (z+\beta)^{-\alpha}$, β large, gives a method essentially equivalent to the typical means of order α of the first kind. M. Riesz has shown that the corresponding abscissa of convergence $\sigma_0(\alpha)$ is a convex function of α. The same is true for the abscissas of uniform and of absolute convergence, $\sigma_u(\alpha)$ and $\sigma_a(\alpha)$. For the Riemann zeta function $\sigma_a(\alpha) = 1-\alpha$, $\alpha \geqq 0$, and $\sigma_u(\alpha) = \frac{1}{2}-\alpha$, $\alpha \geqq \frac{1}{2}$. The interval $0 < \alpha < \frac{1}{2}$ offers all the customary difficulties encountered in the critical strip. (Received November 22, 1933.)

57. Professor D. V. Widder: *Necessary and sufficient conditions for the representation of a function by a doubly infinite Laplace integral.*

We prove in this note that the integral equation $\mu(x) = \int_{-\infty}^{\infty} e^{-xu}\,d\beta(u)$ has an increasing solution $\beta(u)$ if and only if the double integral $\iint \mu(x+y)\,g(x)\,g(y)\,dx\,dy$ is positive (definite or semidefinite). (Received November 15, 1933.)

58. Mr. W. C. Randels: *On the summability of the conjugate series.*

In this note a method of summability is exhibited which is regular and $\tilde{F}$-effective without being $\tilde{L}$-effective. The method is similar to that used by Paley, Rosskopf, and the author in the analogous problem for Fourier series, in a paper shortly to appear in this Bulletin. (Received November 22, 1933.)

59. Mr. M. F. Rosskopf: *Some inequalities for non-uniformly bounded ortho-normal polynomials.* Preliminary report.

An important theorem of Young-Hausdorff-F. Riesz on the Fourier coefficients of a function relative to a set of uniformly bounded ortho-normal polynomials fails to hold in the case of certain types of polynomials which are not

uniformly bounded. The present paper deals with the extension of this theorem to the cases of non-uniformly bounded ortho-normal polynomials such as Legendre, Jacobi, Hermite, Laguerre, and general Tschebycheff polynomials. (Received November 21, 1933.)

60. Professor A. E. Landry: *On the ordered quadrilaterals in- and circumscribed to the plane rational quartic curve (R_2^4) with triangular symmetry.* Preliminary report.

There are two kinds of (ordered) quadrilaterals in- and circumscribed to R_2^4 with triangular symmetry, the one self-symmetric as to a line of symmetry of the curve, the other having no such symmetry. The number, reality, and location of the first kind have been determined by a student of the author and the results published in a doctor's dissertation (June, 1932); the present investigation deals with the same questions for the unsymmetric quadrilaterals. It was found possible to reduce the problem to a final equation in t of the fifth degree. (Received November 27, 1933.)

61. Dr. G. A. Hedlund (National Research Fellow): *Transitive geodesics on closed orientable surfaces of genus greater than one.*

The existence of transitive geodesics on certain closed orientable surfaces of genus greater than one and everywhere non-positive curvature is known. To study the geodesics on such a surface it is desirable to map the covering surface conformally into the unit circle. The existence of transitive geodesics can be proved under the hypothesis that the geodesics and the arcs of circles orthogonal to the unit circle are in one-to-one correspondence in a certain sense. Everywhere non-positive curvature implies this condition, but the converse is not true. The existence of transitive geodesics can then be proved in the presence of regions of positive curvature provided these regions are properly restricted. (Received November 28, 1933.)

62. Professor Philip Franklin: *Geodesics on polyhedral surfaces.*

This paper is concerned with the existence of geodesic lines on a surface of genus zero, having a finite number of conical points. (Received November 28, 1933.)

63. Professor J. L. Synge: *On the deviation of geodesics and null-geodesics particularly in relation to the properties of spaces of constant curvature and indefinite line-element.*

The equation of deviation of geodesics and null-geodesics is obtained very simply. It is applied to give a geometrical interpretation of the Riemannian curvature of a 2-element. For spaces of constant curvature the equation of deviation can be integrated. If the line-element is indefinite, there emanate from a point both convergent and divergent pencils of geodesics, there being convergence only for those directions which make ϵK positive, ϵ being the indicator (± 1) of the pencil (that is, the factor required to make the corresponding

value of the fundamental form positive) and K being the constant curvature. It follows that in space-time with signature $+++-$, of constant positive curvature, all space-like geodesics are closed curves of constant length, but time-like geodesics diverge exponentially. Null-geodesics diverge linearly in a space of constant curvature. (Received November 25, 1933.)

64. Dr. W. W. Flexner: *The intersection of chains on a topological manifold.*

In the present paper the author extends his recent investigations (Annals of Mathematics, (2), vol. 32 (1931)) of topological manifolds to the general case, viz., to intersecting chains C_p and C_q ($s=p+q-n>0$). It is in two parts: (1) the construction on the topological manifold, M_n, of an intersection cycle, Γ_s, using any given covering of M_n by n-cells; (2) the proof that any other cycle, Γ_s', constructed from the same or a different covering is homologous to Γ_s about the geometric intersection of C_p and C_q provided the deformations giving Γ_s and Γ_s' were small enough. (1) Γ_s is constructed piecemeal, using the classical theory on one n-cell of the covering after another. The boundaries of the pieces can be connected by arbitrarily small singular s-chains on M_n giving the cycle. (2) The same sort of construction using chains of a dimension higher gives a chain $C_{s+1}\rightarrow\Gamma_s-\Gamma_s'$. (Received November 8, 1933.)

65. Dr. Deane Montgomery (National Research Fellow): *Operations on plane sets.*

Sierpinski has recently (Mathematica, vol. 5 (1931)) summarized the known results concerning a problem which he formulates as follows: Given a property p of linear point sets and a plane set E, $\Gamma_p(E)$ is the set of all points q on the x-axis such that the vertical line through q cuts E in a set having property p. The problem is to determine the class of $\Gamma_p(E)$ for various classes of sets E and various properties p. The present paper studies as properties p the property of being of the second category, the property of having positive measure, and the property of being at the same time an F_σ and a G_δ. Several different classes of sets E are considered, and an application is made to the theory of implicit functions. (Received November 20, 1933.)

66. Mr. N. E. Steenrod: *Characterizations of certain finite curve-sums.*

The notion of derived aggregates of a closed subset of a compact metric space with respect to a class of closed point sets has been introduced by Whyburn (American Journal of Mathematics, vol. 54, pp. 169–175). He establishes the following: In order that the continuous curve M be expressible as the sum of a finite number of K-curves (where K is some class of continuous curves) it is necessary that, for some integer n, the nth derived aggregate of M with respect to K-curves be vacuous. In the present paper the condition is shown to be sufficient in case K is the class of perfect continuous curves, regular curves, acyclic curves, or boundaries of plane domains. From this it follows that every continuum containing no continuum of condensation and every baum im kleinen curve may be expressed as the sum of two acyclic curves. (Received November 28, 1933.)

67. Professor S. S. Cairns: *Completely regular approximations to regular manifolds.*

For the terms *regular* and *completely regular manifold* we refer to abstract No. 39-5-151, this Bulletin. In the present paper we prove the following approximation theorem. Let M_r be a regular r-manifold in n-space. Then, for any positive values (d, e) however small, there exists a completely regular r-manifold M_r', homeomorphic with M_r in such a way that (1) the distance between any two corresponding points (P, Q) on M_r and M_r' is less than e, and (2) if P is at distance d or more from the irregularities on M_r, then the direction cosines of the tangent r-plane (see *The direction cosines of a p-space in euclidean n-space*, American Mathematical Monthly, vol. 39 (1932), pp. 518–523) to M_r at P differ by less than e from the corresponding direction cosines at Q. The manifold M_r' can be made of class C^m, for any m. With the aid of a polyhedral approximation theorem (abstract No. 39-1-44), the proof is reduced to the case where M_r is a simplicial polyhedron. In that case, M_r' can be made to coincide with M_r at all points of any k-simplex, s_k, which are at distance greater than d from the boundary of $s_k (k=1, \cdots, r)$. (Received December 1, 1933.)

68. Dr. A. H. Black: *Further non-involutorial Cremona space transformations contained in a special linear complex.*

Consider two pencils of surfaces $|F|$ and $|F'|$ of orders n and n', which contain a straight line d as an $(n-1)$- and $(n'-1)$-fold basis element. Make the surfaces of each pencil projective with the points of d. A point P will determine a surface F passing through P, hence a point O on d, and a surface F'. The line PO intersects F' in just one point other than O, the point P' which is the image of P. The general transformation and several special cases are found and discussed. (Received December 1, 1933.)

69. Mr. Garrett Birkhoff: *The topology of transformation-sets.*

The (1, 1) bicontinuous transformations of many spaces, including compact and open manifolds, into themselves, can be connected into continuous groups homeomorphic with subsets of Hilbert space. A study of the explicit content of this fact interrelates closely the convergence properties of abstract spaces and the convergence properties of sets of transformations between them. (Received November 24, 1933.)

70. Mr. Garrett Birkhoff: *Hausdorff groupoids.*

Associated with any Hausdorff space is a topological invariant, namely the Hausdorff space of its $(1\rightarrow 1)$ transformations under a suitable definition of continuity. If the points of the primitive Hausdorff space are isolated, this definition coincides with Kneser's, and we have special applications. Notably, the group of any algebraic extension of an enumerable commutative ring is a Cantor set, metric in the sense of van Dantzig, and possessed of a regular mass-function. Certain related facts concerning groups of importance in algebra are also discussed. (Received November 13, 1933.)

71. Professor Harry Levy: *Linearly connected manifolds and ennuples of curves.*

We give here a treatment of linearly connected manifolds based on ennuples of curves. In place of the coefficients of connection we associate with an ennuple of components ${}_\alpha\lambda^i$ a set of n^3 functions, $\gamma^\alpha_{\beta\gamma}$, invariant under transformations of coordinates. If the ennuple is replaced by a new ennuple of components ${}_\alpha\lambda'^i = l^\beta_\alpha {}_\beta\lambda^i$, the invariants $\gamma^{\alpha}_{\beta\gamma}$ are determined by the equations $l^a_\alpha \gamma^\alpha_{\beta\gamma} = \gamma^a_{bc} l^b_\beta l^c_\gamma + {}_i\lambda^j \partial l^a_\beta / \partial x^j$. On the basis of this law of transformation we are able to develop an absolute calculus very similar to that of Ricci. Among the results we obtain is that one can always select an ennuple whose invariants are zero along a curve, whereas a linearly connected space does not necessarily admit a coordinate system in which the coefficients of connection are zero at a point. We set up functions analogous to the curvature tensor and torsion tensor and discover identities which are generalizations of the well known identities $B^i_{jkl} + B^i_{klj} + B^i_{ljk} = 0$ and $B^i_{jkla} + B^i_{jlak} + B^i_{jakl} = 0$, where B^i_{jkl} and B^i_{jkla} are the Ricci-Christoffel 4 and 5 index symbols of a Riemannian space. Further results are obtained in the case of pseudo-euclidean space, that is, a space of zero curvature but not necessarily of zero torsion. (Received November 28, 1933.)

72. Professor Harry Levy: *Curvatures in Riemannian space.*

For a curve in ordinary space it is well known that $1/\rho = \lim_{P' \to P} \phi/s$ and $1/\tau = \pm \lim_{P' \to P} \theta/s$, where $1/\rho$ is the curvature, $1/\tau$ the torsion, s the arc between P and P', two points of the curve, ϕ the angle between the tangents at P and P', and θ the angle between the binormals. In this note we give an extension of these relations to a curve in Riemannian space. (Received November 28, 1933.)

73. Dr. L. A. Dye and Professor F. R. Sharpe: *The Bertini transformations of space.*

This paper demonstrates the existence of space involutorial transformations such that in each plane of a pencil of invariant planes there is a Bertini transformation. In one type, 6 of the 8 fundamental points of the Bertini transformation lie on a C_6, $p=3$, the other 2 being either fixed or variable on the axis of the pencil of planes. In another type, 6 of the fundamental points lie on a C_6, $p=4$, and the Bertini transformation is degenerate. A third transformation is discussed in which there is a net of invariant quartic surfaces through a C_{11}, $p=14$. The method used in obtaining this last transformation leads also to an involutorial transformation with a net of invariant surfaces of order $n+1$ through a C_{3n-3} of genus $12n-19$. This type has on each plane of a pencil of invariant planes a Geiser transformation having the 7 fundamental points on the C_{3n-3}. (Received November 10, 1933.)

74. Professor J. L. Synge: *Mechanical models of spaces with positive-definite line-element.*

The object of the present paper is to draw attention to the utility of mechanical models as illustrations of spaces defined topologically and metrically.

The configurations of a dynamical system define the points of the manifold and hence its topology; the kinetic energy defines the metric by $ds^2=2T\,dt^2=a_{ij}dx^i dx^j$, this being a positive-definite form. A number of examples are given. (Received November 25, 1933.)

75. Professor B. W. Jones: *A test for equivalence of positive quaternary quadratic forms.*

$f=a_{11}x_1^2+a_{22}x_2^2+a_{33}x_3^2+a_{44}x_4^2+2a_{12}x_1x_2+\cdots$ and f' (similarly expressed) are two positive quaternary quadratic forms of the same Hessian. Four new coefficients are defined by $a_{ii}=-a_{i1}-a_{i2}-a_{i3}-a_{i4}$ $(i=1, 2, 3, 4)$. Unitary transformations are given reducing f to a form ϕ, the sum of whose coefficients $-a_{ij}$ $(i\neq j;\ i, j=1, 2, 3, 4, 5)$ is not more than the sum for any form equivalent to f. No more than two coefficients of ϕ can be positive. Similarly find ϕ'. If no coefficient of ϕ is positive, ϕ' is not equivalent to ϕ unless its coefficients are the coefficients of ϕ in some order. If just one (just two) coefficient of ϕ is positive, ϕ' is not equivalent to ϕ unless just one (just two) of its coefficients is positive, all but four of its coefficients are coefficients of ϕ, *and* the sum of its coefficients is the same as the sum of the coefficients of ϕ. If none of the above tests preclude equivalence, the transformations (which are explicitly stated and few in number) which can take ϕ into ϕ' may be tried. (Received November 25, 1933.)

76. Professor E. V. Huntington: *Independent postulates for an "informal Principia system with equality."*

This paper (which is a continuation of a paper presented in June, 1933, abstract No. 39-7-235) adds to the primitive ideas explicitly mentioned in *Principia Mathematica* the idea of "equal in some undefined respect," and discusses as an illustrative example a system $(K, C, +, ', =)$ in which K is the class of "verdicts" pronounced by a court upon a specified group of defendants; C is the subclass of "correct" verdicts; $+$ is "or"; $'$ is "not"; and $=$ means "equal with respect to legal effect." Some correct verdicts are "truistic," and some incorrect verdicts are "absurd," corresponding to the universe element and the zero element of a Boolean algebra, respectively. A set of independent postulates for such a system is given; and the distinction between the "formal" theory and the "informal" theory in the *Principia* is shown to depend upon the inclusion or rejection of a single postulate. Both these papers will appear in full in this Bulletin. (Received November 14, 1933.)

77. Professor Richard Brauer: *Klein's theory of algebraic equations and its connection with the theory of algebras.*

In his *Vorlesungen über das Ikosaeder*, Klein gave a theory of the equations of the fifth degree. Later he treated the equations of the sixth and seventh degrees in similar manner. In his investigations it appears necessary to adjoin certain accessory irrationalities, which are foreign to the problem. Klein was led to them by special geometric devices, which left the nature of these irrationalities not clear. In this paper it is shown that the question can be treated

in a natural way with the help of the theory of algebras. Thus one obtains an insight into the nature of the accessory irrationalities. (Received December 1, 1933.)

78. Professor C. G. Latimer: *Note on the class number in a rational semi-simple algebra.*

Let $\mathcal{A}$ be a rational semi-simple algebra of order n, and let $\mathcal{G}$ be a domain of integrity of order n in $\mathcal{A}$, employing Dickson's definition. The finiteness of the number of classes of ideals in $\mathcal{G}$ has been shown (a) by Dr. Shover for left ideals in the case where $\mathcal{A}$ is a division algebra (this Bulletin, vol. 39 (1933), pp. 610–614), and (b) by Artin for right ideals in the case where $\mathcal{G}$ is maximal (Mathematisches Seminar, Hamburg, Abhandlungen, vol. 5 (1927), pp. 261–289). In the present note we prove the finiteness of the left ideal class number without the last mentioned restrictions on $\mathcal{A}$ and $\mathcal{G}$. An application is made to similar matrices. By a result due to Dr. Shover (loc. cit.) the right class number is equal to the left class number. (Received December 1, 1933.)

79. Dr. H. W. Raudenbush: *Ideal theory and partial differential equations.*

The object of this paper is to establish an abstract theory of ideals of partial differential forms that will specialize to results obtained by J. F. Ritt. It is proved that every infinite system of forms contains a finite subset such that every form of the system is a root of a linear combination of forms of the subset with forms for coefficients. There follows a decomposition for ideals of forms that contain all derivatives and all roots that are forms. A solution field is constructed and an abstract Hilbert-Netto theorem is obtained for forms. (Received November 29, 1933.)

80. Dr. D. H. Lehmer: *Lacunary recurrences for Bernoulli numbers.*

Recent work on Fermat's last theorem has made desirable an extension of the tables of Bernoulli numbers. This paper gives a number of explicit recurrence formulas for B_n which are particularly designed to facilitate the calculation of these large numbers. In contrast to the recurrences used heretofore, in which B_n was made to depend on each preceding B, the present recurrences have gaps so that B_n may be computed from those preceding B's whose subscripts are congruent to n with respect to some modulus m. The kth term of the recurrence, then, has the form $B_{n-km}\alpha(n, m, k)c_k$, in which $\alpha(n, m, k)$ is a certain binomial coefficient, and where $c_1, c_2, c_3, \cdots$, is itself a recurring series of fixed order with constant coefficients, and depends only on m. As m is increased the recurrence becomes shorter, while the c's become more complicated. The case of $m=12$ gives the most practical formula. The method of deriving these formulas is quite elementary and makes use of the Blissard-Lucas umbral calculus. Similar recurrences are obtained for Euler's numbers and the coefficients of the tangent function. The method may also be applied to the coefficients of certain elliptic functions. (Received November 29, 1933.)

81. Professor Marie J. Weiss: *On simply transitive primitive groups.*

Certain properties of the transitive constituents of the subgroup that fixes one letter of a simply transitive primitive group are analyzed in this paper. The chief results are embodied in the following theorems: If the subgroup that fixes one letter of a simply transitive primitive group has two transitive constituents of relatively prime degrees m and n, $n>m$, it has a transitive constituent of degree $(>n)$ a divisor of mn. If the subgroup that fixes one letter of a simply transitive primitive group has a regular constituent of degree pq, where p and q are primes, it is of order pq. (Received November 27, 1933.)

82. Mr. M. M. Flood: *Division by non-singular matric polynomials.*

It is shown in this paper how all divisions of a matric polynomial P by a non-singular matric polynomial A may be obtained. Wedderburn has shown the existence of the division transformation in this case, and a simpler proof of this is included here. Some restrictions on the degree and on the coefficients of the quotients are found. The matric polynomial V is said to "reduce" A if VA is of lesser degree than A. The matric polynomial W is said to be "associated" with A if the product WA is of the same degree as A and has leading coefficient unity. All polynomials associated with A are easily found and their degree is unique. Obviously all polynomials which reduce A are determined. (Received November 27, 1933.)

83. Dr. L. M. Blumenthal (National Research Fellow): *A determinant theorem obtained from the characterization of pseudo r-spheric sets.*

The characterization of pseudo d-cyclic sets of points given by the author in two recent papers (American Journal of Mathematics, vol. 54 (1932), pp. 729–738; vol. 55 (1933), pp. 619–640) leads to a geometric proof of the following determinant theorem: *If* $\Delta=|r_{ij}|$, $r_{ij}=r_{ji}$ $(i, j=1, 2, \cdots, n)$, $r_{ii}=1$, *of order* $n>4$, *is such that* (i) *the absolute value of* r_{ij}, $i\neq j$, *is less than* 1, (ii) *every third-order principal minor vanishes,* (iii) *at least one fourth-order principal minor does not vanish, then* (a) *no fourth-order principal minor vanishes,* (b) *each* r_{ij}, $i\neq j$, *has the absolute value* $1/2$, *and* (c) *the value of the determinant is* $-(1/2)\cdot(3/2)^{n-1}(n-3)$. It is suggested that this theorem is the first link of a chain of theorems on determinants of the form Δ of order $n>k+3$, the kth link of which contains the following hypotheses: (i) every principal minor of order less than $k+2$ is positive, (ii) every principal minor of order $k+2$ vanishes, (iii) at least one principal minor of order $k+3$ does not vanish; the conclusions being (a) no principal minor of order $k+3$ vanishes, (b) each element r_{ij}, $i\neq j$, has absolute value $1/(k+1)$, and (c) the value of the determinant is $-[1/(k+1)]\cdot[(k+2)/(k+1)]^{n-1}(n-k-2)$. These theorems do not seem to be in the literature of determinant theory, and no purely algebraic method of deducing them is known to the author. (Received November 8, 1933.)

84. Dr. A. T. Craig: *Correlation of indices.*

If the variables x, y, z, with mean values zero, are normally correlated, it is proved that the indices $u=x/y$, $v=z/y$ are correlated in accord with a Cauchy function with linear regression and with a coefficient of correlation equal to the partial correlation coefficient of x and z. Extensions are made to more than two indices and to the case in which the components of the indices are non-normally correlated but possess linear regression systems. (Received November 25, 1933.)

85. Dr. A. T. Craig: *Note on the moments of a Bernoulli distribution.*

Romanovsky (Biometrika, vol. 15 (1923), pp. 410–412) has given a recursion formula for the moments of the Bernoulli distribution $(q+p)^n$ about its mean np. In the present note, a simpler method of arriving at this formula is given. The method is further used to find a recursion formula for the moments of the Poisson exponential. (Received November 25, 1933.)

86. Dr. Saunders MacLane: *Problems of structure-theoretic type in mathematical logic.*

An analysis of the classical form of mathematical logic shows that this logic is concerned chiefly with the study of the detailed structure of mathematical theorems. This suggests several new types of problems, such as the one of finding standard methods of "translating" ordinary mathematical language into logic. More generally, logic may consider the structure, not only of a single theorem, but also of any body of mathematical doctrine, such as a proof or a theory. In particular, the investigation of the structure of proofs leads on the one hand to an analysis of many "processes" of proof, and on the other hand to the study of purposes of proof. By means of a number of examples, it is shown that such purposes can be objectively analysed into completely general "plans" and "methods." (Received November 25, 1933.)

87. Dr. Saunders MacLane: *General properties of algebraic systems.*

Any algebraic variety, such as a group, a field, a group with operators, and the like, concerns essentially a system composed of a number of functions. If this notion of a system be extended to include relations as well, then we can say that any abstract mathematical theory consists of some system and a number of axioms and theorems about this system. The object of this paper is the investigation of the properties of systems. In the first place, many typical algebraic concepts, such as isomorphism, homeomorphism, direct product, abstraction with respect to a congruence, etc., are really relations between systems and can be defined with complete generality. From this standpoint, we can prove a number of interesting theorems concerning the interconnections of these relations and including as special cases many well known theorems of algebra. One such theorem is the generalization to systems of the "second isomorphism theorem" for groups. Many algebraic theorems may be viewed

as special cases of the general theorem that any two isomorphic systems have the same structure. This theorem, as well as a number of similar general theorems, are proved by an application of certain logical concepts; some interesting connections between algebra and mathematical logic are thereby indicated. (Received November 25, 1933.)

88. Professor A. A. Albert: *On a certain algebra of quantum mechanics.*

P. Jordan, J. von Neumann, and E. Wigner have considered certain commutative, non-associative algebras of quantum mechanics. Their algebras have finite order over the field of all real numbers, are real in the sense of Artin, and obey the law $(xy)x^2=x(yx^2)$. In particular, any algebra M of real n-rowed square matrices and with multiplication in M defined as quasi-multiplication $ab=\frac{1}{2}(a\cdot b+b\cdot a)$ of matrices a, b in M satisfies their postulates. Moreover they have proved that every irreducible algebra satisfying their properties is either an algebra M or is the algebra N of all three-rowed Hermitian matrices with elements in the real Cayley algebra of order eight, multiplication being again defined as quasi-multiplication of the matrices. In the present paper I prove that this latter algebra is a new algebra, not equivalent to any M above. Moreover I give a brief proof of the complicated property $(xy)x^2=x(yx^2)$. (Received November 29, 1933.)

89. Professor A. A. Albert: *On cyclic equations of prime degree.*

If $F(x)$ is cyclic of odd prime degree p over F and e is a primitive pth root of unity, then $F(x, e)=F(y, e)$, $y^p=c(e)=c$ in $K=F(e)$, $c\neq d^p$ for any d of K. (Compare Fricke's *Algebra*, I, pp. 413–415.) But a false converse is given in the literature. In fact I prove that necessarily $c(e^t)=d^pc^t$, where d is in K and K is cyclic of degree n over F with generating automorphism $e\longrightarrow e^t$. But then conversely if $y^p=c$ as above, the field $F(y)$ is cyclic of degree pn over F and $F(y)=F(e)\times F(x)$, where $F(x)$ is cyclic of degree p over F. I also show how to construct certain quantities c giving all cyclic $F(x)$ so that I have in fact given a formal construction of all cyclic fields of odd prime degree. The results are used to prove that a necessary and sufficient condition that a normal division algebra D of degree p over F be cyclic is that D shall contain a sub-field $F(z)$, $z^p=g$ in F. (Received November 29, 1933.)

90. Dr. R. D. James (National Research Fellow): *The value of the constants in Waring's problem.*

The constants which appear in the Hardy-Littlewood analysis for the number $G(k)$ in Waring's problem were evaluated in a previous paper (abstract No. 39-3-103, this Bulletin). It was then possible to deduce new results for $g(k)$. In another paper (abstract No. 39-1-26, this Bulletin) values for $G(k)$ when k is odd were obtained which were an improvement on those obtained by the Hardy-Littlewood method. It is the purpose of the present paper to evaluate the constants of the second paper and so obtain new results for $g(k)$ when k is odd. It is shown, for example, that $g(7)\leqq 281$, $g(9)\leqq 1023$; that is, *all*

numbers are sums of at most 281 seventh powers and of at most 1023 ninth powers. The previous results were $g(7) \leqq 353$ and $g(9) \leqq 1252$. (Received November 29, 1933.)

91. Professor C. H. Forsyth: *An interpolation formula which when adjusted for rational centering of data proves to be osculatory.*

At the Atlantic City meeting of the Society in 1932, the author offered certain modifications for several known interpolation formulas which had the effect of treating given equi-spaced data as areas instead of as values concentrated at the midpoints of the successive intervals. In the present paper it is shown that such modifications for probably the most important interpolation formula (Newton's) prove to be osculatory, that is, each pair of partial interpolation curves for two successive intervals of interpolation have common osculating circles, and therefore the same curvature (including slope) at their point of intersection. Hence, the whole series of interpolations over any number of successive intervals of interpolation will prove relatively smooth from interval to interval. Incidentally, this formula can be used to only third differences and still retain the osculatory property. Heretofore, osculatory interpolation formulas were thought to require fifth differences. (Received November 29, 1933.)

92. Dr. H. L. Krall: *On some asymptotic relations for the characteristic values of elliptic differential equations.*

Let $L(u)=Au_{xx}+2Bu_{xy}+Cu_{yy}+Du_x+Eu_y+Fu$, where $AC-B^2>0$ and the coefficients have continuous partial derivatives of the first three orders. Also let $M(u)$ be the adjoint operator. An asymptotic relation is given for the characteristic values of the system $L(u)=\lambda v$, $M(v)=\lambda u$ with the boundary conditions $u=0$, $v=0$. A condition is also found for the characteristic values of the system $L(u)=\lambda u$ with $u=0$ on the boundary. These results are generalizations of the results of Geppert and Gheorghiu who discussed the case $L(u)=u_{xx}+u_{yy}+Du_x+Eu_y+Fu$. (Received December 2, 1933.)

93. Professor R. E. Langer: *The asymptotic solutions of certain linear ordinary differential equations of the second order.*

The ordinary differential equation $du^2/dz^2+\lambda p_1(z,\lambda)(du/dz)+\lambda^2 p_2(z,\lambda)=0$ in which λ is a complex parameter and the functions $p_i(z,\lambda)$ are expansible in descending powers of λ includes as special cases many equations of importance. A change of variable gives it the form $d^2u/dz^2-\{\lambda^2 q_0(z)+\lambda q_1(z)+q_2(z,\lambda)\}u=0$ in which q_2 is a series in λ^{-1}. If z is real and on an interval in which the determinations of $\{q_0(z)\}^{1/2}$ are distinct, the asymptotic forms of the solutions have been known. This paper considers the equation with the variable z complex and in a region within which $q_0(z)$ has a zero of the second order. The asymptotic forms of the solutions are derived. The paper supplements earlier ones by the author on equations of similar type in which, however, $q_1(z)$ was taken to be identically zero. (Received December 8, 1933.)

94. Mr. E. P. Northrop: *Note on a singular integral.*

This note consists of two theorems regarding convergence in the mean of order 2, as $m \to \infty$, of the integral $(2\pi)^{-1/2}\int_{-\infty}^{+\infty} f(u)K(x-u; m)du$, where $f(u)$ is an arbitrary function belonging to $L_2(-\infty, +\infty)$, and $K(u; m)$ belongs to $L_2(-\infty, +\infty)$ for every m. It is found that a set of necessary and sufficient conditions for the convergence of the integral to $f(x)$ [to zero] is as follows: (i) The Fourier transform of $K(u; m)$ is less than or equal to a constant M for all m and almost all u. (ii) The Fourier transform of $K(u; m)$ converges in the mean of order 2 to 1 [to zero] over every finite interval. (Received December 8, 1933).

95. Professor J. L. Walsh: *Note on the orthogonality of Tchebycheff polynomials on confocal ellipses.*

This note shows that the Tchebycheff polynomials $p_n(z)$, orthogonal on the interval $(-1 \leqq z \leqq +1)$ with respect to the norm function $(1-z^2)^{-1/2}$, are orthogonal on every ellipse C whose foci are the points -1 and $+1$ with respect to the norm function $n(z) = |1-z^2|^{-1/2}$, in the sense $\int_C n(z)p_n(z)\overline{p_m(z)}\,|dz| = 0$, $m \neq n$. (Received December 15, 1933.)

96. Professor I. A. Barnett and Dr. D. S. Nathan: (National Research Fellow): *Linear transformations in function space.*

The authors establish the group property for the one-parameter family of linear transformations in function-space generated by a given infinitesimal linear integral transformation in this space. They accomplish this without employing computational methods, such as were used by Kennison (National Academy Proceedings, vol. 16, pp. 607–609). They then extend the result to a composite function-space. They also extend to a composite function-space Kowalewski's results on the orthogonal group in function-space (Wiener Sitzungsberichte, vol. 120, II, pp. 101–109). In particular, they obtain the orthogonal group in both finite and infinitesimal form; they show that a given infinitesimal orthogonal transformation generates a one-parameter group of orthogonal transformations, and they obtain the analogue of the Cayley formulas for expressing the coefficients of an orthogonal transformation in terms of skew-symmetric quantities. (Received December 27, 1933.)

97. Dr. Leo Zippin: *Semi-Peanian spaces.*

We investigate in Cauchy spaces (complete metric separable) the notion of semi-compactness: a topologic space is semi-compact if *for every point* x and any U_x (neighborhood of x) there exists a $V_x \subset U_x$ such that the boundary, $F(V_x)$, is compact (or vacuous). We show first that a semi-compact Cauchy space can be compactified by the "addition" of a countable infinity of points. If the space is moreover connected and locally connected we call it semi-Peanian (=quasi-Peanian+semi-compact). These spaces are *characterized* by the following property: a semi-Peanian space C can always be compactified to a Peanian space C^* such that $C^* - C \equiv H$ is a totally disconnected F_σ-set which is *thoroughly avoidable* in C^*. This means that if D is any open connected subset

of C^* then $D-D\cdot H$ is connected. It is interesting that C^* is an invariant of C. As consequences of the theorem above, we have an extension to semi-Peanian spaces of most of the theory of Peanian spaces, including the recent work of Claytor on planar Peanian spaces. (Received December 27, 1933.)

98. Dr. E. R. van Kampen: *Topological characterizations of 2-dimensional manifolds.*

Characterizations are given of the 2-sphere, 2-cell with and without boundary, 2-dimensional manifolds without boundary, infinite 2-dimensional manifolds and regions of the sphere. Only those for the two types of general manifolds are new. The others are to be found in two papers by Zippin in the American Journal of Mathematics, vols. 52 and 55, but here the proofs are considerably simplified. The wording of the different characterizations, as well as the proofs, is similar to a very large extent. (Received December 27, 1933.)

99. Professor P. A. Smith: *A theorem on fixed points.*

Let S be a subspace of a cartesian S_m and such that every continuous image in S of every hypersphere of dimension $\leqq pm-m-1$ can be deformed in S to a point; then every topological transformation of S into itself which is of period p leaves fixed at least one point. (Received December 19, 1933.)

100. Dr. Charles Hopkins: *Metabelian groups of order p^m.*

The discussion is restricted chiefly to non-divisible metabelian groups G of order p^m. These are classified primarily according to the number n of independent generating operations and secondarily according to the structure of the commutator subgroup. For $p>2$ it is proved that a basis $s_1, s_2, \cdots, s_n$ always exists such that $s_1^{\lambda_n} s_2^{\lambda_2} \cdots s_n^{\lambda_1}=1$ only when each λ is divisible by the order of the corresponding s. The existence of certain categories of groups G is demonstrated by exhibiting a method of construction. Necessary and sufficient conditions that a correspondence $s_i \sim \prod_{j=1}^{n} s_j^{\lambda_{ij}}$, $i=1, \cdots, n$, define an automorphism of G are developed. A method for constructing every automorphism of G is given. Several interesting applications of this theory will be presented in a later paper. (Received December 13, 1933.)

101. Dr. G. D. Gore: *On generalized inscribed sequences of surfaces.*

The theory of conjugate nets and of asymptotic families of curves on surfaces was generalized by Bompiani (Rendiconti del Circolo Matematico di Palermo, vol. 46 (1922), p. 91). Generalizations of the classical sequences of Laplace were developed from Bompiani's results by B. Segre (Annales Scientifiques de L'École Normale Supérieure, vol. 44 (1927), p. 152). The present paper generalizes the classical inscribed sequence of Laplace and discovers a large class of sequences more general than those of Segre. Four types of the generalized inscribed sequences are inscribed in the sequences of Segre. The results show that the transformations of Laplace and Segre are applicable to a much less restricted class of surfaces than the class to which they have been applied. (Received December 13, 1933.)

102. Professor W. E. Roth: *On Zehfuss matrices and applications.*

If $A=(a_{ij})$, $(i=1, 2, \cdots, m; j=1, 2, \cdots, n)$, is an $m\times n$ matrix and B is a $p\times q$ matrix, then the $mp\times nq$ matrix $(a_{ij}B)$, whose elements occur in mn blocks of p rows and q columns, each of which is the corresponding element of B multiplied by a_{ij}, is a Zehfuss matrix of A and B (Zehfuss, Zeitschrift für Mathematik und Physik, vol. 3 (1858), p. 298). (Rutherford recently discussed the conditions of equality of such matrices (this Bulletin, vol. 39 (1933), pp. 801–808)). The transformation to certain normal forms and the characteristic values of Zehfuss matrices are studied; also their elementary divisors in certain special cases are taken up when it is assumed that those of $A-\lambda I$ and of $B-\mu I$ are known. Application is then made to the solution of the linear matrix equation $A_1\times B_1+A_2\times B_2+\cdots+A_r\times B_r=C$, and to the solution of the following problem: To write down explicitly the equation $F(x)=0$ of degree mn whose roots are $y_{ij}=\phi(\alpha_i; \beta_j)$ $(i=1, 2, \cdots, m; j=1, 2, \cdots, n)$, where $\phi(\lambda, \mu)$ is a given polynomial in λ and μ and α_i and β_j are the roots of the algebraic equations $f(x)=0$ and $g(x)=0$ respectively. (Received December 13, 1933.)

103. Mr. K. W. Wegner: *Equivalence of pairs of Hermitian matrices.*

Two pairs of Hermitian matrices A, B and C, D are said to be equivalent if and only if there exists a non-singular matrix T such that $\overline{T}'AT=C$ and $\overline{T}'BT=D$. In the case in which $|B|\neq 0$ and $|D|\neq 0$, the necessary and sufficient conditions for the equivalence of A, B and C, D are: (1) the elementary divisors of $A-\lambda B$ and $C-\lambda D$ are the same; (2) the indices of $B(B^{-1}A-\lambda I)^n$ and $D(D^{-1}C-\lambda I)^n$ are the same for all positive integral n and real λ. Condition (1) is not sufficient as has sometimes been stated. The method used is much simpler than that used by Muth (Journal für die reine und angewandte Mathematik, 128 (1905), 302–21) in obtaining the conditions for the real equivalence of pairs of real symmetric matrices. It is based on the most general matrix commutative with the classical canonical form of $B^{-1}A$ and $D^{-1}C$. The conditions are easily generalized to the non-singular case in which neither of the determinants $|\rho A+\sigma B|$ and $|\rho C+\sigma D|$ is zero identically in ρ and σ. The singular case is under consideration. (Received December 8, 1933.)

NEW PUBLICATIONS

PART I. PURE MATHEMATICS

ANDERSON (W. E.). A first course in college mathematics. New York, Harper, 1933. 10+326 pp.+92 pp.

BEETLE (R. D.). McGraw-Hill five-place logarithmic and trigonometric tables. New York, McGraw-Hill, 1933. 8+132 pp.

BERNSTEIN (V.). Leçons sur les progrès récents de la théorie des séries de Dirichlet. Avec une préface de J. Hadamard. (Collection Borel.) Paris, Gauthier-Villars, 1933. 16+320 pp.

BOEHM (C.). See EULER (L.).

BOHDE (H.). Mathematisch-technische Tafeln. 7te, vermehrte Auflage. Berlin, Springer, 1933. 105 pp.

BOLL (M.). See CARNAP (R.).

BRADLEY (A. D.). The geometry of repeating design and geometry of design for high schools. New York, Teachers College, 1933. 6+132 pp.

BRITISH ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE. Mathematical tables. Volume 3: Minimum decompositions into fifth powers. Prepared by L. E. Dickson. (Published under the supervision of the British Association Committee for the Calculation of Mathematical Tables.) London, British Association, 1933. 6+368 pp.

BROWN (O. E.). See MORRIS (M.).

CARNAP (R.). L'ancienne et la nouvelle logique. Traduction du Général E. Vouillemin, introduction de Marcel Boll. (Actualités Scientifiques et Industrielles, No. 76.) Paris, Hermann, 1933. 36 pp.

CARTAN (E.). Sur la structure des groupes de transformations finis et continus. Paris, Vuibert, 1933.

DATTA (B.). The science of the Sulba. A study in early Hindu geometry. (Readership lectures for the year 1931.) Calcutta, University of Calcutta, 1932. 16+240 pp.

DAVIS (H. T.), edited by. Tables of the higher mathematical functions. Volume 1. (Published as a contribution of the Waterman Institute for Scientific Research, Indiana University.) Bloomington, Ind., Principia Press, 1933. 377+13 pp.

DICKSON (L. E.). See BRITISH ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE.

DINGLER (H.). Die Grundlagen der Geometrie. Ihre Bedeutung für Philosophie, Mathematik, Physik und Technik. Stuttgart, Enke, 1933. 84 pp.

DURELL (C. V.) and ROBSON (A.). Elementary calculus. Volume 1. London, G. Bell, 1933. 8+240 pp.

EULER (L.). Opera omnia. Series I: Opera mathematica. Volumen XVI, sectio I: Commentationes analyticae ad theoriam serierum infinitarum pertinentes. Edidit C. Boehm. Leipzig, Teubner, 1933. 10+355 pp.

EWING (C. A.). Plane trigonometry. New York, McGraw-Hill, 1933. 12+166 pp.

FAVARD (J.). Leçons sur les fonctions presque-périodiques. Préface de G. Julia. (Cahiers Scientifiques.) Paris, Gauthier-Villars, 1933. 8+183 pp.

GERARD (A. E.). See THOMPSON (S. P.).

GIBSON (G. A.) and PINKERTON (P.). Elements of analytical geometry. Chapters I–VI: The straight line. London, Macmillan, 1933. 8+105+6 pp.

GLIEBE (J. J.). The mathematical atom: its involution and evolution exemplified in the trisection of the angle; a problem in plane geometry. 3d edition, revised. San Francisco, St. Boniface Franciscan Friary, and London, Technical Records, 1933. 3+87 pp.

GODEAUX (L.). Questions non résolus de géométrie algébrique. (Actualités Scientifiques et Industrielles, No. 77: Exposés sur l'analyse mathématique publiés sous la direction de J. Hadamard, I.) Paris, Hermann, 1933. 24 pp.

GÜNTHER (G.). Grundzüge einer neuen Theorie des Denkens in Hegels Logik. Leipzig, Meiner, 1933. 226 pp.

HADAMARD (J.). See BERNSTEIN (V.), GODEAUX (L.).

HAHN (H.). Logik, Mathematik und Naturerkennen. Wien und Leipzig, Carl Gerold's Sohn, 1933. 33 pp.

HART (W. L.). Plane trigonometry. New York, Heath, 1933. 6+186+16 pp.; tables, 4+124 pp.

HEIDEL (W. A.). The heroic age of science. The conception, ideals and methods of science among the ancient Greeks. Baltimore, Williams and Wilkins, 1933. 203 pp.

INGE (D.). God and the astronomers. Containing the Warburton Lectures, 1931–1933. London and New York, Longmans, 1933. 13+308 pp.

JULIA (G.). See FAVARD (J.).

KOWALEWSKI (G.). Lehrbuch der höheren Mathematik. Band 3: Fortsetzung der Differential- und Integralrechnung. Differentialgleichungen. Differentialgeometrie. Funktionen einer komplexen Veränderlichen. Probleme der Variationsrechnung. Berlin, de Gruyter, 1933. 2+252 pp.

LIETZMANN (W.). Kegelschnittlehre. Leipzig, Teubner, 1933. 46 pp.

LINDOW (M.). Gewöhnliche Differentialgleichungen. 2te Auflage. Leipzig, Teubner, 1933. 121 pp.

——— Integralrechnung. 4te Auflage. Leipzig, Teubner, 1933. 102 pp.

MACE (C. A.). The principles of logic. An introductory survey. London and New York, Longmans, 1933. 13+388 pp.

MORRIS (M.) and BROWN (O. E.). Differential equations. New York, Prentice-Hall, 1933. 14+410 pp.

PINKERTON (P.). See GIBSON (G. A.).

PRICE (E. A.). Logical sequence in plane geometry. Rapid revision by means of ideograms. Cambridge, University Press, 1933. 12 pp.

RAMACHANDRAN (G. V.). See RANGANATHAN (V.).

RANGANATHAN (V.) and RAMACHANDRAN (G. V.). Intermediate trigonometry. Trichinopoly, Wednesday Review Press, 1933. 225 pp.

ROBSON (A.). See DURELL (C. V.).

ROSENBACH (J. B.) and WHITMAN (E. A.). College algebra. Boston, Ginn, 1933. 12+394 pp.

SAKS (S.). Théorie de l'intégrale. (Monografje Matematyczne, II.) Warsaw, Mathematical Seminar of the University of Warsaw, 1933. 7+290 pp.

SILBERSTEIN (L.). Causality a law of nature or a maxim of the naturalist. London, Macmillan, 1933. 159 pp.

THOMPSON (S. P.). Le calcul intégral et différentiel à la portée de tout le monde. Traduit de l'anglais par A. E. Gerard. 3e édition, revue et corrigée. Paris, Dunod, 1933. 8+290 pp.

VÉRONNET (A.). Le calcul vectoriel. Cours d'algèbre de mathématiques spéciales et de mathématiques générales. Préface par H. Villat. Paris, Gauthier-Villars, 1933. 18+252 pp.

VILLAT (H.). See VÉRONNET (A.).

VOUILLEMIN (E.). See CARNAP (R.).

WAREHAM (A. L.). The foundations of the universe. London, C. W. Daniel, 1933. 43 pp.

WHITMAN (E. A.). See ROSENBACH (J. B.).

WITTING (A.). Integralrechnung. (Sammlung Göschen.) Berlin, de Gruyter, 1933. 175 pp.

PART II. APPLIED MATHEMATICS

ALTER (D.). Introduction to practical astronomy. New York, Crowell, 1933. 8+80 pp.

APPELL (P.). Traité de mécanique rationnelle. 2e édition. Tome 5. Revue par R. Thiry. Paris, Gauthier-Villars, 1933. 198 pp.

ASTBURY (W. T.). Fundamentals of fibre structure. London, Oxford University Press, 1933. 10+187 pp.

BLEICH (F.). Stahlhochbauten, ihre Theorie, Berechnung und bauliche Gestaltung. Band 2. Berlin, Springer, 1933. 5+376 pp.

BORCHARDT (W. G.). First course in mechanics. London, Rivingtons, 1933. 238 pp.

——— A second course in mechanics. London, Rivingtons, 1933.

BRÖTZ (O.). See SCHARRER (G.).

BROWN (E. W.) and SHOOK (C. A.). Planetary theory. New York, Macmillan, and Cambridge, University Press, 1933. 12+302 pp.

CALDERWOOD (J. P.). See MOYER (J. A.).

CROWDER (W. F.). See DAVIES (G. R.).

DAVIES (G. R.) and CROWDER (W. F.). Methods of statistical analysis in the social sciences. New York, Wiley, and London, Chapman and Hall, 1933. 11+355 pp.

DENNISON (B. C.). See KARAPETOFF (V.).

EDWARDS (H. W.). Analytic and vector mechanics. (International Series in Physics.) New York and London, McGraw-Hill, 1933. 10+428 pp.

EVANS (R. D.). See WAGNER (C. F.).

GEIGER (H.) und SCHEEL (K.), herausgegeben von. Handbuch der Physik. 2te Auflage. Band 24, Teil 1: Quantentheorie. Redigiert von A. Smekal. Berlin, Springer, 1933. 9+853 pp.

GEMANT (A.). Liquid dielectrics. English translation by V. Karapetoff. (National Research Council Committee on Electrical Insulation, Monograph No. 2.) New York, Wiley, and London, Chapman and Hall, 1933. 9+185 pp.

GERLACH (WALTHER) und GERLACH (WERNER). Die chemische Emissions-Spektralanalyse. Teil 2. Leipzig, Voss, 1933. 191 pp.

GERLACH (WERNER). See GERLACH (WALTHER).

GIBSON (C. R.). Electrical conceptions of today. A lucid explanation of many of the latest theories concerning atoms, electrons and other matters relating to electricity. London, Seeley, Service and Company, 1933. 284 pp.

GOLDING (E. W.). Electrical measurements and measuring instruments. (Engineering Degree Series.) London, Pitman, 1933. 804 pp.

GRIMSEHL (E.). A textbook of physics. Edited by R. Tomaschek. Authorized translation from the 7th German edition by L. A. Woodward. Volume 3: Electricity and magnetism. London and Glasgow, Blackie, 1933. 14+685 pp.

GRODZINKSI (P.) und POLSTER (H.). Getriebelehre. Band 1: Geometrische Grundlagen. Band 2: Angewandte Getriebelehre. (Sammlung Göschen.) Berlin, de Gruyter, 1933. 136+142 pp.

GUERNER (J.). See SCHWAIGER (A.).

GUGGENHEIM (E. A.). Modern thermodynamics by the methods of Willard Gibbs. London, Methuen, 1933. 16+206 pp.

GUSTAFSON (T.). Über den Magnuseffekt nach der asymptotischen hydrodynamischen Theorie. (Dissertation, Lund.) Lund, Håkan Ohlssons Buchdruckerei, 1933.

HARBURGER (W.). Strahlender Raum. München, Duncker und Humbolt, 1933. 95 pp.

HOHENEMSER (K.) und PRAGER (W.). Dynamik der Stabwerke. Eine Schwingungslehre für Bauingenieure. Berlin, Springer, 1933. 6+367 pp.

HOWARD (H. B.). The stresses in aeroplane structures. London, Pitman, 1933. 274 pp.

INFELD (L.). Nowe drogi nauki: kwanty i materja. (Z Dziedziny Nauki i Techniki, Tom 2.) Warsaw, Mathesis Polskiej, 1933. 10+284 pp.

KARAPETOFF (V.). Experimental electrical engineering and manual for electrical testing. 4th edition, completely revised and reset. Volume 1. Revised by B. C. Dennison. New York, Wiley, and London, Chapman and Hall, 1933. 28+781 pp.

——— See GEMANT (A.).

KING (H. W.) and WISLER (C. O.). Hydraulics. 3d edition, revised. New York, Wiley, and London, Chapman and Hall, 1933. 12+292 pp.

KLEIN (P. E.). Transformatoren und Drosseln. Berlin-Tempelhof, J. Schneider, 1933. 83 pp.

KUCZYNSKI (R. R.). Fertility and reproduction. Methods of measuring the balance of births and deaths. New York, Falcon Press, 1932. 3+94 pp.

LENSE (J.). Reihenentwicklung in der mathematischen Physik. Berlin, de Gruyter, 1933. 178 pp.

LERTES (P.). Elektrische Musik. Leipzig, Steinkopff, 1933. 11+207 pp.

LOWE (E. A.). Direct and alternating currents. Theory and machinery. New York and London, McGraw-Hill, 1933. 13+656 pp.

MASSEY (H. S. W.). See MOTT (N. F.).

MOELWYN-HUGHES (E. A.). The kinetics of reactions in solution. Oxford, Clarendon Press, and London, Oxford University Press, 1933. 7+313 pp.

MORECROFT (J. H.). Electron tubes and their application. New York, Wiley, and London, Chapman and Hall, 1933. 13+458 pp.

MOTT (N. F.) and MASSEY (H. S. W.). The theory of atomic collisions. (International Series of Monographs on Physics.) Oxford, Clarendon Press, and New York, Oxford University Press, 1933. 15+283 pp.

MOYER (J. A.), CALDERWOOD (J. P.), and POTTER (A. A.). Elements of engineering thermodynamics. 5th edition, rewritten and reset. New York, Wiley, and London, Chapman and Hall, 1933. 14+192 pp.

NEMÉNYI (P.). Wasserbauliche Strömungslehre. Leipzig, Barth, 1933. 275 pp.

PLASKETT (H. H.). The place of observation in astronomy. New York, Oxford University Press, 1933. 32 pp.

POLSTER (H.). See GRODZINKSI (P.).

PONCIN (H.). Recherches sur le mouvement d'un fluide pesant dans un plan vertical. (Publications Scientifiques du Ministère de l'Air, No. 16.) Paris, Gauthier-Villars, 1933. 108 pp.

———— Sur les cavitations de forme permanente. Préface de H. Villat. (Publications Scientifiques du Ministère de l'Air, No. 18.) Paris, Gauthier-Villars, 1933. 126 pp.

POTTER (A. A.). See MOYER (J. A.).

PRAGER (W.). See HOHENEMSER (K.).

QUITTNER-BERTOLASI (E.). Das Verhältnis von Trend und Konjunkturzyklen als mathematisch-ökonomisches Problem. (Veröffentlichung der Frankfurter Gesellschaft für Konjunkturforschung.) Leipzig, Hans Buske Verlag, 1933. 57 pp.

RAPSON (E. T. A.). Electrical transmission and distribution. London, Oxford University Press, 1933. 179 pp.

RESTA (O.). La derivazione anomola nel tiro delle artiglierie. Bologna, Azzoguidi, 1933.

ROEVER (W. H.). The Mongean method of descriptive geometry according to the procedure of Gino Loria. New York, Macmillan, 1933. 14+151 pp.

RUTHERFORD (E.). The artificial transmutation of the elements. Being the thirty-fifth Robert Boyle Lecture, delivered before the Oxford University Junior Scientific Club on June 2, 1933. London, Oxford University Press, 1933. 12 pp.

SALIGER (R.). Der Eisenbeton, seine Berechnung und Gestaltung. 6te ergänzte Auflage. Leipzig, Verlag Kröner, 1933. 661 pp.

SCHARRER (G.) und BRÖTZ (O.). Gebäudeschwingungen. (Forschungsheft 359.) Berlin, V. D. I. Verlag, 1933. 24 pp.

SCHEEL (K.). See GEIGER (H.).

SCHÖLER (R.). Die Statik und Festigkeitslehre einschliesslich der Theorie, Berechnungen und Konstruktionen des Eisenbetonbaues. Band 1. Leipzig, Verlag Voigt, 1933. 175 pp.

SCHWAIGER (A.). Calcul pratique des lignes de transport d'énergie électrique: basse, moyenne, haute tension et lignes à longue distance. Traduction et adaptation d'après l'édition allemande par J. Guerner. Paris, Dunod, 1933. 12+142 pp.

SHOOK (C. A.). See BROWN (E. W.).

SMEKAL (A.). See GEIGER (H.).

THIRY (R.). See APPELL (P.).

TOMASCHEK (R.). See GRIMSEHL (E.).

UNOLD (G.). Die praktische Berechnung der Stahlskelettrahmen. Berlin, Wilhelm Ernst, 1933. 9+75 pp.

VILLAT (H.). See PONCIN (H.).

WAGNER (C. F.) and EVANS (R. D.). Symmetrical components as applied to the analysis of unbalanced electrical circuits. New York and London, McGraw-Hill, 1933. 16+437 pp.

WARBURG (E.). Lehrbuch der Experimentalphysik für Studierende. 23te und 24te verbesserte Auflage. Dresden und Leipzig, Steinkopf, 1933. 484 pp.

WISLER (C. O.). See KING (H. W.).

WOLF (R.). Geschichte der Astronomie. (Reprint of the 1877 edition.) München, Oldenbourg, und Leipzig, Koehlers Antiquarium, 1933. 16+815 pp.

WOODWARD (L. A.). See GRIMSEHL (E.).

WULF (T.). Die Faden-Elektrometer. Berlin, Dümmlers Verlag, 1933. 147 pp.

OFFICIAL COMMUNICATIONS

Meetings of the Society have been fixed at the following times and places:

The usual February Meeting in New York City will be omitted.

NEW YORK CITY, March 30-31, 1934.

Abstracts must be in the hands of Associate Secretary J. R. Kline, 501 West 116th St., New York City, not later than March 9. Abstracts received by the Secretaries by that date will appear in the March issue of the BULLETIN. By invitation of the program committee, there will be a symposium on *Algebraic geometry*, under the leadership of Professors A. B. Coble, Solomon Lefschetz, Oscar Zariski, and Virgil Snyder.

CHICAGO, ILLINOIS, April 6-7, 1934.

Abstracts must be in the hands of Associate Secretary M. H. Ingraham, University of Wisconsin, Madison, Wis., not later than March 9. Abstracts received by the Secretaries by that date will appear in the March issue of the BULLETIN. At this meeting, Professor R. E. Langer will give an address on *The asymptotic solution of linear ordinary differential equations*.

BERKELEY, CALIFORNIA, June 20, 1934.

Abstracts must be in the hands of Associate Secretary T. M. Putnam, University of California, Berkeley, Calif., not later than May 30. Abstracts received by the Secretaries by May 6 will appear in the May issue of the BULLETIN.

R. G. D. RICHARDSON, *Secretary of the Society.*

Articles for insertion in the BULLETIN should be addressed to E. R. HEDRICK, Editor of the BULLETIN, University of California at Los Angeles. Reviews should be sent to W. R. LONGLEY, Yale University, New Haven, Conn. Notes should be sent to H. W. KUHN, Ohio State University, Columbus, Ohio.

Subscriptions to the BULLETIN, orders for back numbers, and inquiries in regard to non-delivery of current numbers should be addressed to the American Mathematical Society, 450-459 Ahnaip St., Menasha, Wis., or 501 West 116th St., New York.

The initiation fees and the annual dues of members of the Society (see this BULLETIN, p. 322, May, 1930; and the List of Officers and Members, September, 1932, p. 66), are payable to the Treasurer of the Society, Professor G. W. Mullins, 501 West 116th St., New York City.

WHOLE NUMBER 412

CONTENTS

PAGE

The Thanksgiving Meeting in Pasadena. By T. M. Putnam 1

The Thanksgiving Meeting in Cincinnati. By M. H. Ingraham 4

Birkhoff on Aesthetic Measure. By C. A. Garabedian ... 7

Pappus's Collection. By David Eugene Smith 11

Banach on Linear Operations. By J. D. Tamarkin 13

Saks on Integration. By J. D. Tamarkin 16

Shorter Notices 19

Notes 25

Abstracts of Papers 28

New Publications 59

For official Communications and Notices, see the inside of the back cover.

GEORGE BANTA PUBLISHING COMPANY, MENASHA, WISCONSIN